

Chapter 1

GWA in ωk space

1.1 Green function $G^{(0)}$

$$G^{(0)}(\zeta_1, \zeta_2; \omega) = \sum_i \frac{\phi_i^{(0)}(\zeta_1)\phi_i^{(0)*}(\zeta_2)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)}$$

$$G^{(0)}(r_1, r_2; \omega) = \sum_i \frac{\phi_i^{(0)}(r_1)\phi_i^{(0)*}(r_2)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)}$$

$$\begin{aligned} G^{(0)}(r_1, r_2; \omega) &= \sum_{\xi_1 \xi_2} G^{(0)}(\zeta_1, \zeta_2; \omega) \\ &= \sum_{\xi_1 \xi_2} \sum_i \frac{\phi_i^{(0)}(\zeta_1)\phi_i^{(0)*}(\zeta_2)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \\ &= \sum_{\xi_1} \sum_i \frac{\phi_i^{(0)}(\zeta_1)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \left(\phi_i^{(0)*}(r_2, \xi_2 = -1/2) + \phi_i^{(0)*}(r_2, \xi_2 = +1/2) \right) \\ &= \sum_{\xi_1} \sum_i \frac{\phi_i^{(0)}(\zeta_1)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \phi_i^{(0)*}(r_2) \\ &= \sum_i \left(\phi_i^{(0)}(r_1, \xi_1 = -1/2) + \phi_i^{(0)}(r_1, \xi_1 = +1/2) \right) \frac{\phi_i^{(0)*}(r_2)}{\omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \end{aligned}$$

1.2 Screened Interaction

$$W(\zeta_1, \zeta_2, \omega) = \int d\zeta_3 \varepsilon^{-1}(\zeta_1, \zeta_3, \omega) w(\zeta_3, \zeta_2)$$

$$W(\zeta_1, \zeta_2, \omega) = W(r_1, r_2, \omega) = \int dr_3 \varepsilon^{-1}(r_1, r_3, \omega) w(r_3, r_2) \quad \text{spin independent (but it's not the case of } \varepsilon)$$

$$w(r_1, r_2) = \frac{1}{|r_1 - r_2|} \quad \text{coulombian many body interaction}$$

$$w(q, G) = \frac{4\pi}{|q + G|^2} \quad \text{fourier transform of the coulombian interaction}$$

$$W(r_1, r_2, \omega) = \int dr_3 \varepsilon^{-1}(r_1, r_3, \omega) w(r_3, r_2)$$

$$\begin{aligned} W(r_1, r_2, \omega) &= \int dr_3 \frac{1}{(2\pi)^3} \int_{\text{BZ}} dq \sum_{G_1, G_3} e^{i(q+G_1)r_1} \varepsilon^{-1}(q, G_1, G_3, \omega) e^{-i(q+G_3)r_3} . \\ &\quad \cdot \frac{1}{(2\pi)^3} \int_{\text{BZ}} dq_1 \sum_{G_2} e^{i(q_1+G_2)r_3} \frac{4\pi}{(q_1 + G_2)^2} e^{-i(q_1+G_2)r_2} \\ &\quad \int dr_3 e^{-i(q+G_3)r_3} e^{i(q_1+G_2)r_3} = (2\pi)^3 \delta(q - q_1) \delta(G_3 - G_2) \end{aligned}$$

$$W(r_1, r_2, \omega) = \frac{1}{(2\pi)^3} \int_{\text{BZ}} dq \sum_{G_1, G_2} e^{i(q+G_1)r_1} \varepsilon^{-1}(q, G_1, G_2, \omega) \frac{4\pi}{(q + G_2)^2} e^{-i(q+G_2)r_2}$$

$$\begin{aligned} W(r_1, r_2, \omega) &= \frac{1}{\Omega} \sum_{q, G_1, G_2} e^{i(q+G_1)r_1} W(q, G_1, G_2, \omega) e^{-i(q+G_2)r_2} \\ &= \frac{1}{\Omega} \sum_{q, G_1, G_2} e^{i(q+G_1)r_1} \varepsilon^{-1}(q, G_1, G_2, \omega) \frac{4\pi}{|q + G_2|^2} e^{-i(q+G_2)r_2} \end{aligned}$$

$$W(q, G_1, G_2, \omega) = \varepsilon^{-1}(q, G_1, G_2, \omega) \frac{4\pi}{|q + G_2|^2} = \bar{\varepsilon}^{-1}(q, G_1, G_2, \omega) \frac{4\pi}{|q + G_1||q + G_2|}$$

$\bar{\varepsilon}$ = symmetrized epsilon.

1.3 Single Plasmon Pole Model

$$\varepsilon^{-1}(\omega) = \delta + \frac{\Omega^2}{\omega^2 - \tilde{\omega}^2}$$

$$\varepsilon^{-1}(G, G', q, \omega) = \delta(G, G') + \frac{\Omega^2(G, G', q)}{\omega^2 - \tilde{\omega}^2(G, G', q)}$$

$$A(G, G', q) = -\frac{\Omega^2(G, G', q)}{\tilde{\omega}^2(G, G', q)} = \varepsilon^{-1}(G, G', q, 0) - \delta(G, G')$$

$$\tilde{\omega}^2(G, G', q) = \left[\frac{A(G, G', q)}{\varepsilon^{-1}(G, G', q, 0) - \varepsilon^{-1}(G, G', q, iE_0)} - 1 \right] E_0^2$$

$$\varepsilon^{-1}(iE_0) = \delta + \frac{\Omega^2}{-E_0^2 - \tilde{\omega}^2}$$

$$\varepsilon^{-1}(iE_0)[E_0^2 + \tilde{\omega}^2] = \delta[E_0^2 + \tilde{\omega}^2] - \frac{\Omega^2}{\tilde{\omega}^2} \tilde{\omega}^2 = \delta E_0^2 + \varepsilon^{-1}(0) \tilde{\omega}^2$$

$$\tilde{\omega}^2 = \frac{\varepsilon^{-1}(iE_0) - \delta}{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)} E_0^2$$

$$= \left[\frac{\varepsilon^{-1}(iE_0) - \delta}{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)} + \frac{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)}{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)} - 1 \right] E_0^2$$

$$\tilde{\omega}^2 = \left[\frac{\varepsilon^{-1}(0) - \delta}{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)} - 1 \right] E_0^2 = \left[\frac{A}{\varepsilon^{-1}(0) - \varepsilon^{-1}(iE_0)} - 1 \right] E_0^2$$

1.4 GW approximation for the Self-Energy

$$\tilde{\Sigma}_M^{GW}(x_1, x_2) = iG(x_1, x_2)W(x_1^+, x_2)$$

$$\tilde{\Sigma}_M^{GW}(\zeta_1, \zeta_2, \omega) = \frac{i}{2\pi} \int d\omega' e^{-i\omega'\eta} G(\zeta_1, \zeta_2, \omega - \omega') W(\zeta_1, \zeta_2, \omega')$$

W is spin-independent and G does not take factors 2 on spin sum:

$$\tilde{\Sigma}_M^{GW}(r_1, r_2, \omega) = \frac{i}{2\pi} \int d\omega' e^{-i\omega'\eta} G(r_1, r_2, \omega - \omega') W(r_1, r_2, \omega')$$

$$\begin{aligned} \tilde{\Sigma}_M^{GW}(r_1, r_2, \omega) &= \frac{i}{2\pi} \int d\omega' e^{-i\omega'\eta} \sum_i \frac{\phi_i^{(0)}(r_1)\phi_i^{(0)*}(r_2)}{\omega - \omega' - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \cdot \\ &\quad \cdot \frac{1}{\Omega} \sum_{q, G_1, G_2} e^{i(q+G_1)r_1} \bar{\varepsilon}^{-1}(q, G_1, G_2, \omega') \frac{4\pi}{|q + G_1||q + G_2|} e^{-i(q+G_2)r_2} \end{aligned}$$

1.5 Matrix elements of the Self-Energy operator

$$\begin{aligned} \langle \phi_j^{(0)}(r_1) | \tilde{\Sigma}_M^{GW}(r_1, r_2, \omega) | \phi_j^{(0)}(r_2) \rangle &= \frac{i}{2\pi} \frac{1}{\Omega} \sum_{q, G_1, G_2} \frac{4\pi}{|q + G_1||q + G_2|} \sum_i \cdot \\ &\quad \cdot \int dr_1 \phi_j^{(0)*}(r_1) e^{i(q+G_1)r_1} \phi_i^{(0)}(r_1) \int dr_2 \phi_i^{(0)*}(r_2) e^{-i(q+G_2)r_2} \phi_j^{(0)}(r_2) \cdot \\ &\quad \cdot \int_{-\infty}^{+\infty} d\omega' e^{-i\omega'\eta} \frac{1}{\omega - \omega' - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \bar{\varepsilon}^{-1}(q, G_1, G_2, \omega') \end{aligned}$$

$$i = \{n_i, k_i\}$$

$$\rho_{ij}(G) = \int dr \phi_i^{(0)*}(r) e^{-i(q+G)r} \phi_j^{(0)}(r) \quad \text{with} \quad q = k_j - k_i - G_0, \quad q \in 1\text{BZ}$$

$$\begin{aligned} \langle j | \tilde{\Sigma}_M^{GW}(\omega) | j \rangle &= \frac{i}{2\pi} \frac{1}{\Omega} \sum_i \sum_{G_1, G_2} \delta(q - (k_j - k_i - G_0)) \frac{4\pi}{|q + G_1||q + G_2|} \rho_{ij}^*(G_1) \rho_{ij}(G_2) \cdot \\ &\quad \cdot \int_{-\infty}^{+\infty} d\omega'' e^{i\omega''\eta} \frac{1}{\omega'' + \omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \bar{\varepsilon}^{-1}(q, G_1, G_2, \omega'') \end{aligned}$$

$$\begin{aligned} \varepsilon^{-1}(q, G, G', \omega) &\rightarrow \delta(G, G') \rightarrow \Sigma_x \quad \text{Sigma exchange} \\ \varepsilon^{-1}(q, G, G', \omega) &\rightarrow \frac{\Omega^2(q, G, G')}{\omega^2 - \tilde{\omega}^2(q, G, G')} \rightarrow \Sigma_c \quad \text{Sigma correlation} \end{aligned}$$

1.6 Self-Energy: exchange term

$$\begin{aligned} \langle j | \tilde{\Sigma}_x^{\text{GW}}(\omega) | j \rangle &= \frac{i}{2\pi} \frac{1}{\Omega} \sum_i \sum_{G_1} \delta(q - (k_j - k_i - G_0)) \frac{4\pi}{|q + G_1|^2} |\rho_{ij}(G_1)|^2 \cdot \\ &\quad \cdot \oint_{\text{upper half plane}} d\omega'' e^{i\omega''\eta} \frac{1}{\omega'' + \omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \end{aligned}$$

Only the poles of G in the upper half imaginary ω -plane are included in the closed, anti-clockwise path. They correspond to the particle poles (excitation corresponding to occupied states). The integral yields ($f_i = 0, 1$ occupation number):

$$\oint d\omega'' \dots = 2\pi i f_i \operatorname{Res}_i = 2\pi i f_i$$

$$\langle j | \tilde{\Sigma}_x^{\text{GW}} | j \rangle = -\frac{4\pi}{\Omega} \sum_i f_i \sum_{G_1} \delta(q - (k_j - k_i - G_0)) \frac{1}{|q + G_1|^2} |\rho_{ij}(G_1)|^2$$

it does not depend on ω because it constitutes only a shift term of the poles along the real axis which doesn't change the integral.

1.7 Self-Energy: correlation term

$$\begin{aligned} \langle j | \tilde{\Sigma}_c^{\text{GW}}(\omega) | j \rangle &= \frac{i}{2\pi} \frac{1}{\Omega} \sum_i \sum_{G_1, G_2} \delta(q - (k_j - k_i - G_0)) \frac{4\pi}{|q + G_1||q + G_2|} \rho_{ij}^*(G_1) \rho_{ij}(G_2) \cdot \\ &\quad \cdot \oint_{\text{uhp-acw}} d\omega'' e^{i\omega''\eta} \frac{1}{\omega'' + \omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \frac{\Omega^2(q, G_1, G_2)}{\omega''^2 - (\tilde{\omega}(q, G_1, G_2) - i\eta)^2} \\ &\quad \oint_{\text{uhp-acw}} d\omega'' e^{i\omega''\eta} \frac{1}{\omega'' + \omega - \epsilon_i^{(0)} + i\eta \operatorname{sgn}(\epsilon_i^{(0)} - \mu)} \frac{1}{(\omega'' - \tilde{\omega} + i\eta)(\omega'' + \tilde{\omega} - i\eta)} \end{aligned}$$

The included poles are in $\omega'' = -\tilde{\omega} + i\eta$ and if $f_i = 1$ in $\omega'' = \epsilon_i^{(0)} - \omega + i\eta$.

$$\begin{aligned} \oint_{\text{uhp-acw}} d\omega'' \dots &= 2\pi i \left(\frac{f_i}{(\epsilon_i^{(0)} - \omega)^2 - \tilde{\omega}^2} + \frac{1}{\omega - \tilde{\omega} - \epsilon_i^{(0)}} \frac{1}{-2\tilde{\omega}} \right) \\ &= -i\pi \left(\frac{1}{\tilde{\omega}(\omega - \tilde{\omega} - \epsilon_i^{(0)})} - \frac{2f_i}{(\omega - \epsilon_i^{(0)})^2 - \tilde{\omega}^2} \right) \\ &= -i\pi \left(\frac{\omega - \epsilon_i^{(0)} + \tilde{\omega}}{\tilde{\omega}(\omega - \epsilon_i^{(0)} + \tilde{\omega})(\omega - \epsilon_i^{(0)} - \tilde{\omega})} - \frac{\tilde{\omega}2f_i}{\tilde{\omega}(\omega - \epsilon_i^{(0)} + \tilde{\omega})(\omega - \epsilon_i^{(0)} - \tilde{\omega})} \right) \\ &= -i\pi \frac{\omega - \epsilon_i^{(0)} - \tilde{\omega}(2f_i - 1)}{\tilde{\omega}(\omega - \epsilon_i^{(0)} + \tilde{\omega})(\omega - \epsilon_i^{(0)} - \tilde{\omega})} \frac{\omega - \epsilon_i^{(0)} + \tilde{\omega}(2f_i - 1)}{\omega - \epsilon_i^{(0)} + \tilde{\omega}(2f_i - 1)} \\ &= -i\pi \frac{(\omega - \epsilon_i^{(0)})^2 - \tilde{\omega}^2(2f_i - 1)^2}{(\omega - \epsilon_i^{(0)})^2 - \tilde{\omega}^2} \frac{1}{\tilde{\omega}(\omega - \epsilon_i^{(0)} + \tilde{\omega}(2f_i - 1))} \\ &= -i\pi \frac{1}{\tilde{\omega}(\omega - \epsilon_i^{(0)} + \tilde{\omega}(2f_i - 1))} \end{aligned}$$

as $(2f_i - 1)^2$ is always 1.

$$\begin{aligned} \langle j | \tilde{\Sigma}_c^{\text{GW}}(\omega) | j \rangle &= \frac{2\pi}{\Omega} \sum_i \sum_{G_1, G_2} \delta(q - (k_j - k_i - G_0)) \frac{\rho_{ij}^*(G_1) \rho_{ij}(G_2)}{|q + G_1||q + G_2|} \cdot \\ &\quad \cdot \frac{\Omega^2(q, G_1, G_2)}{\tilde{\omega}(q, G_1, G_2)(\omega - \epsilon_i^{(0)} + \tilde{\omega}(q, G_1, G_2)(2f_i - 1))} \end{aligned}$$

Appendix A

Definitions, Fourier transforms

A.1 Definitions, notations

$$\Omega = N_k \Omega_{\text{cell}} \quad \text{Crystal Volume}$$

$$\Omega_{\text{BZ}} = \frac{(2\pi)^3}{\Omega_{\text{cell}}}$$

$$\frac{1}{(2\pi)^3} \int_{\text{BZ}} dk = \frac{1}{\Omega} \sum_k^{\text{BZ}}$$

$$\frac{1}{\Omega_{\text{BZ}}} \int_{\text{BZ}} dk = \frac{1}{N_k} \sum_k^{\text{BZ}}$$

A.2 Fourier transform definition

$$f(\omega) = \int dt e^{-i\omega t} f(t) \quad \text{direct fourier transform}$$

$$f(t) = \frac{1}{2\pi} \int d\omega f(\omega) e^{i\omega t} \quad \text{inverse fourier transform}$$

A.3 Fourier transform of a two lattice indices quantity

$$\begin{aligned}
f(q, G_1, G_2) &= \frac{1}{(2\pi)^3} \int dr_1 dr_2 e^{-i(q+G_1)r_1} f(r_1, r_2) e^{i(q+G_2)r_2} \\
f(r_1, r_2) &= \frac{1}{(2\pi)^3} \int_{\text{BZ}} dq \sum_{G_1, G_2} e^{i(q+G_1)r_1} f(q, G_1, G_2) e^{-i(q+G_2)r_2} \\
f(r_1, r_2) &= \frac{1}{\Omega} \sum_{q, G_1, G_2} e^{i(q+G_1)r_1} f(q, G_1, G_2) e^{-i(q+G_2)r_2}
\end{aligned}$$

Demonstration:

$$\begin{aligned}
f(Q_1, Q_2) &= \frac{1}{(2\pi)^3} \int dr_1 dr_2 e^{-iQ_1 r_1} f(r_1, r_2) e^{iQ_2 r_2} \quad \text{definition fourier transform} \\
f(r_1, r_2) &= \frac{1}{(2\pi)^3} \int dQ_1 dQ_2 e^{iQ_1 r_1} f(Q_1, Q_2) e^{-iQ_2 r_2} \quad \text{definition inverse fourier transform} \\
f(r_1 + R, r_2 + R) &= f(r_1, r_2) \quad \text{lattice periodicity} \\
\frac{1}{(2\pi)^3} \int dQ_1 dQ_2 e^{iQ_1(r_1+R)} f(Q_1, Q_2) e^{-iQ_2(r_2+R)} &= \frac{1}{(2\pi)^3} \int dQ_1 dQ_2 e^{iQ_1 r_1} f(Q_1, Q_2) e^{-iQ_2 r_2} \\
\frac{1}{(2\pi)^3} \int dq_1 dq_2 \sum_{G_1, G_2} e^{i(q_1+G_1)(r_1+R)} f(q_1, q_2, G_1, G_2) e^{-i(q_2+G_2)(r_2+R)} &= \\
&= \frac{1}{(2\pi)^3} \int dq_1 dq_2 \sum_{G_1, G_2} e^{i(q_1+G_1)r_1} f(q_1, q_2, G_1, G_2) e^{-i(q_2+G_2)r_2} \\
e^{iG_1 R} &= e^{-iG_2 R} = 1 \\
e^{i(q_1-q_2)R} &= 1 \quad \Rightarrow \quad q_1 = q_2 \quad \Rightarrow \quad f(q_1, q_2, G_1, G_2) = f(q_1, G_1, G_2) \delta(q_1 - q_2)
\end{aligned}$$

A.4 Case $q \rightarrow 0, G = 0$ for $\rho^2(q, G = 0)/q^2$

$$F = \frac{1}{\Omega} \sum_q^{\text{BZ}} \frac{f(q)}{q^2} = \frac{1}{\Omega} f(q=0) I_{\text{SZ}} + \frac{1}{\Omega} \sum_{q \neq 0}^{\text{BZ}} \frac{f(q)}{q^2}$$

$$I_{\text{SZ}} = \frac{\Omega}{(2\pi)^3} \int_{\Omega_{\text{BZ}}/N_k} d\mathbf{q} \frac{1}{q^2} = \frac{N_k}{\Omega_{\text{BZ}}} \int_{\Omega_{\text{BZ}}/N_k} d\mathbf{q} \frac{1}{q^2}$$

If we assume a spheric Brillouin zone of volume V and radius $(3V/4\pi)^{1/3}$:

$$\frac{1}{V} \int_V d\mathbf{q} \frac{1}{q^2} = \frac{4\pi}{V} \int_0^{(3V/4\pi)^{1/3}} dq = 3^{1/3} (4\pi)^{2/3} V^{-2/3}$$

$$I_{\text{SZ}} = 7.79 \left(\frac{\Omega_{\text{BZ}}}{N_k} \right)^{-2/3}$$

In the case of a Brillouin Zone shape such for an fcc material:

$$I_{\text{SZ}} = 7.44 \left(\frac{\Omega_{\text{BZ}}}{N_k} \right)^{-2/3}$$