

The Isabelle/HOL Algebra Library

Clemens Ballarin
Florian Kammüller
Lawrence C Paulson

November 22, 2007


Contents

1	Orders and Lattices	6
1.1	Partial Orders	6
1.1.1	Upper	7
1.1.2	Lower	7
1.1.3	least	8
1.1.4	greatest	8
1.2	Lattices	9
1.2.1	Supremum	9
1.2.2	Infimum	10
1.3	Total Orders	12
1.4	Complete lattices	12
1.5	Examples	14
1.5.1	Powerset of a Set is a Complete Lattice	14
2	Monoids and Groups	14
2.1	Definitions	14
2.2	Cancellation Laws and Basic Properties	18
2.3	Subgroups	19
2.4	Direct Products	20
2.5	Homomorphisms and Isomorphisms	21
2.6	Commutative Structures	22
2.7	The Lattice of Subgroups of a Group	24
3	Product Operator for Commutative Monoids	24
3.1	Inductive Definition of a Relation for Products over Sets	24
3.2	Left-Commutative Operations	25
3.3	Commutative Monoids	27
3.4	Products over Finite Sets	28

4	The Combinatorial Argument Underlying the First Sylow Theorem	30
4.1	Prime Theorems	30
4.2	Exponent Theorems	31
4.3	Main Combinatorial Argument	32
5	Cosets and Quotient Groups	33
5.1	Basic Properties of Cosets	34
5.2	Normal subgroups	36
5.3	More Properties of Cosets	37
5.3.1	Set of Inverses of an r -coset.	38
5.3.2	Theorems for $\langle \# \rangle$ with $\# \rangle$ or $\langle \#$	38
5.3.3	An Equivalence Relation	38
5.3.4	Two Distinct Right Cosets are Disjoint	39
5.4	Further lemmas for r -congruent	39
5.5	Order of a Group and Lagrange's Theorem	39
5.6	Quotient Groups: Factorization of a Group	40
5.7	The First Isomorphism Theorem	41
6	Sylow's Theorem	43
6.1	Main Part of the Proof	43
6.2	Discharging the Assumptions of <i>syLOW-central</i>	44
6.2.1	Introduction and Destruct Rules for H	44
6.3	Equal Cardinalities of M and the Set of Cosets	45
6.3.1	The Opposite Injection	46
6.4	Sylow's Theorem	47
7	Bijections of a Set, Permutation Groups and Automorphism Groups	47
7.1	Bijections Form a Group	48
7.2	Automorphisms Form a Group	48
8	Abelian Groups	49
8.1	Basic Properties	49
8.2	Sums over Finite Sets	52
9	The Algebraic Hierarchy of Rings	54
9.1	Basic Definitions	54
9.2	Rings	54
9.2.1	Normaliser for Rings	55
9.2.2	Sums over Finite Sets	57
9.3	Integral Domains	57
9.4	Fields	58
9.5	Morphisms	58

10 Modules over an Abelian Group	60
10.1 Definitions	60
10.2 Basic Properties of Algebras	61
11 Univariate Polynomials	62
11.1 The Constructor for Univariate Polynomials	62
11.2 Effect of Operations on Coefficients	63
11.3 Polynomials Form a Commutative Ring.	64
11.4 Polynomials Form an Algebra	66
11.5 Further Lemmas Involving Monomials	67
11.6 The Degree Function	68
11.7 Polynomials over Integral Domains	70
11.8 The Evaluation Homomorphism and Universal Property	71
11.9 Sample Application of Evaluation Homomorphism	73
12 More Lifting from Groups to Abelian Groups	75
12.1 Definitions	75
12.2 Cosets	76
12.3 Subgroups	78
12.4 Normal additive subgroups	78
12.4.1 Definition of <i>abelian-subgroup</i>	78
12.5 Congruence Relation	81
12.6 Factorization	82
12.7 The First Isomorphism Theorem	83
12.8 Homomorphisms	83
13 Lemmas Lifted from CosetExt.thy	85
13.1 General Lemmas from AlgebraExt.thy	85
13.2 Lemmas for Right Cosets	85
13.3 Lemmas for the Set of Right Cosets	86
13.4 Addition of Subgroups	86
14 Ideals	87
14.1 General definition	87
14.2 Ideals Generated by a Subset of <i>carrier R</i>	87
14.3 Principal Ideals	87
14.4 Maximal Ideals	88
14.5 Prime Ideals	88
15 Properties of Ideals	89
15.1 Special Ideals	89
15.2 General Ideal Properties	89
15.3 Intersection of Ideals	89
15.3.1 Intersection of a Set of Ideals	89

15.4	Addition of Ideals	90
15.5	Ideals generated by a subset of <i>carrier</i> R	90
15.5.1	Generation of Ideals in General Rings	90
15.5.2	Generation of Principal Ideals in Commutative Rings	91
15.6	Union of Ideals	92
15.7	Properties of Principal Ideals	92
15.8	Prime Ideals	92
15.9	Maximal Ideals	93
15.10	Derived Theorems Involving Ideals	93
16	Homomorphisms of Non-Commutative Rings	94
16.1	The kernel of a ring homomorphism	95
16.2	Cosets	95
17	QuotRing: Quotient Rings	96
17.1	Multiplication on Cosets	96
17.2	Quotient Ring Definition	96
17.3	Factorization over General Ideals	96
17.4	Factorization over Prime Ideals	97
17.5	Factorization over Maximal Ideals	97
18	The Ring of Integers	98
18.1	Some properties of <i>int</i>	98
18.2	The Set of Integers as Algebraic Structure	98
18.2.1	Definition of \mathcal{Z}	98
18.2.2	Interpretations	98
18.2.3	Generated Ideals of \mathcal{Z}	100
18.2.4	Ideals and Divisibility	100
18.2.5	Ideals and the Modulus	100
18.2.6	Factorization	101



theory *Lattice* **imports** *Main* **begin**

1 Orders and Lattices

Object with a carrier set.

```
record 'a partial-object =
  carrier :: 'a set
```

1.1 Partial Orders

```
record 'a order = 'a partial-object +
  le :: ['a, 'a] => bool (infixl  $\sqsubseteq_1$  50)
```

```
locale partial-order =
  fixes L (structure)
  assumes refl [intro, simp]:
     $x \in \text{carrier } L \implies x \sqsubseteq x$ 
  and anti-sym [intro]:
     $\llbracket x \sqsubseteq y; y \sqsubseteq x; x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x = y$ 
  and trans [trans]:
     $\llbracket x \sqsubseteq y; y \sqsubseteq z; x \in \text{carrier } L; y \in \text{carrier } L; z \in \text{carrier } L \rrbracket \implies x \sqsubseteq z$ 
```

```
constdefs (structure L)
  less :: [-, 'a, 'a] => bool (infixl  $\sqsubset_1$  50)
   $x \sqsubset y \iff x \sqsubseteq y \ \& \ x \not\sim y$ 
```

— Upper and lower bounds of a set.

```
Upper :: [-, 'a set] => 'a set
Upper L A == {u. (ALL x. x ∈ A ∩ carrier L --> x ⊆ u)} ∩
  carrier L
```

```
Lower :: [-, 'a set] => 'a set
Lower L A == {l. (ALL x. x ∈ A ∩ carrier L --> l ⊆ x)} ∩
  carrier L
```

— Least and greatest, as predicate.

```
least :: [-, 'a, 'a set] => bool
least L l A == A ⊆ carrier L & l ∈ A & (ALL x : A. l ⊆ x)
```

```
greatest :: [-, 'a, 'a set] => bool
greatest L g A == A ⊆ carrier L & g ∈ A & (ALL x : A. x ⊆ g)
```

— Supremum and infimum

```
sup :: [-, 'a set] => 'a ( $\sqcup_1$ -[90] 90)
 $\sqcup A == \text{THE } x. \text{least } L \ x \ (\text{Upper } L \ A)$ 
```

$inf :: [-, 'a\ set] ==> 'a\ (\sqcap_1\ [90]\ 90)$
 $\sqcap A ==\ THE\ x.\ greatest\ L\ x\ (Lower\ L\ A)$

$join :: [-, 'a, 'a] ==> 'a\ (\mathbf{infixl}\ \sqcup_1\ 65)$
 $x\ \sqcup\ y ==\ sup\ L\ \{x, y\}$

$meet :: [-, 'a, 'a] ==> 'a\ (\mathbf{infixl}\ \sqcap_1\ 70)$
 $x\ \sqcap\ y ==\ inf\ L\ \{x, y\}$

1.1.1 Upper

lemma *Upper-closed* [*intro, simp*]:

$Upper\ L\ A \subseteq carrier\ L$
 $\langle proof \rangle$

lemma *UpperD* [*dest*]:

fixes L (**structure**)
shows $[| u \in Upper\ L\ A; x \in A; A \subseteq carrier\ L |] ==> x \sqsubseteq u$
 $\langle proof \rangle$

lemma *Upper-memI*:

fixes L (**structure**)
shows $[| !! y. y \in A ==> y \sqsubseteq x; x \in carrier\ L |] ==> x \in Upper\ L\ A$
 $\langle proof \rangle$

lemma *Upper-antimono*:

$A \subseteq B ==> Upper\ L\ B \subseteq Upper\ L\ A$
 $\langle proof \rangle$

1.1.2 Lower

lemma *Lower-closed* [*intro, simp*]:

$Lower\ L\ A \subseteq carrier\ L$
 $\langle proof \rangle$

lemma *LowerD* [*dest*]:

fixes L (**structure**)
shows $[| l \in Lower\ L\ A; x \in A; A \subseteq carrier\ L |] ==> l \sqsubseteq x$
 $\langle proof \rangle$

lemma *Lower-memI*:

fixes L (**structure**)
shows $[| !! y. y \in A ==> x \sqsubseteq y; x \in carrier\ L |] ==> x \in Lower\ L\ A$
 $\langle proof \rangle$

lemma *Lower-antimono*:

$A \subseteq B ==> Lower\ L\ B \subseteq Lower\ L\ A$
 $\langle proof \rangle$

1.1.3 least

lemma *least-carrier* [*intro, simp*]:
shows $\text{least } L \ l \ A \ ==> \ l \in \text{carrier } L$
<proof>

lemma *least-mem*:
 $\text{least } L \ l \ A \ ==> \ l \in A$
<proof>

lemma (*in partial-order*) *least-unique*:
 $[\text{least } L \ x \ A; \text{least } L \ y \ A] \ ==> \ x = y$
<proof>

lemma *least-le*:
fixes L (**structure**)
shows $[\text{least } L \ x \ A; a \in A] \ ==> \ x \sqsubseteq a$
<proof>

lemma *least-UpperI*:
fixes L (**structure**)
assumes *above*: $!! \ x. \ x \in A \ ==> \ x \sqsubseteq s$
and *below*: $!! \ y. \ y \in \text{Upper } L \ A \ ==> \ s \sqsubseteq y$
and $L: A \subseteq \text{carrier } L \ s \in \text{carrier } L$
shows $\text{least } L \ s \ (\text{Upper } L \ A)$
<proof>

1.1.4 greatest

lemma *greatest-carrier* [*intro, simp*]:
shows $\text{greatest } L \ l \ A \ ==> \ l \in \text{carrier } L$
<proof>

lemma *greatest-mem*:
 $\text{greatest } L \ l \ A \ ==> \ l \in A$
<proof>

lemma (*in partial-order*) *greatest-unique*:
 $[\text{greatest } L \ x \ A; \text{greatest } L \ y \ A] \ ==> \ x = y$
<proof>

lemma *greatest-le*:
fixes L (**structure**)
shows $[\text{greatest } L \ x \ A; a \in A] \ ==> \ a \sqsubseteq x$
<proof>

lemma *greatest-LowerI*:
fixes L (**structure**)
assumes *below*: $!! \ x. \ x \in A \ ==> \ i \sqsubseteq x$
and *above*: $!! \ y. \ y \in \text{Lower } L \ A \ ==> \ y \sqsubseteq i$

and $L: A \subseteq \text{carrier } L \ i \in \text{carrier } L$
shows *greatest* $L \ i$ (*Lower* $L \ A$)
 $\langle \text{proof} \rangle$

1.2 Lattices

locale *lattice* = *partial-order* +
assumes *sup-of-two-exists*:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies \exists s. \text{least } L \ s \ (\text{Upper } L \ \{x, y\})$
and *inf-of-two-exists*:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies \exists s. \text{greatest } L \ s \ (\text{Lower } L \ \{x, y\})$

lemma *least-Upper-above*:
fixes L (**structure**)
shows $\llbracket \text{least } L \ s \ (\text{Upper } L \ A); x \in A; A \subseteq \text{carrier } L \rrbracket \implies x \sqsubseteq s$
 $\langle \text{proof} \rangle$

lemma *greatest-Lower-above*:
fixes L (**structure**)
shows $\llbracket \text{greatest } L \ i \ (\text{Lower } L \ A); x \in A; A \subseteq \text{carrier } L \rrbracket \implies i \sqsubseteq x$
 $\langle \text{proof} \rangle$

1.2.1 Supremum

lemma (**in** *lattice*) *joinI*:
 $\llbracket \exists l. \text{least } L \ l \ (\text{Upper } L \ \{x, y\}) \rrbracket \implies P \ l; x \in \text{carrier } L; y \in \text{carrier } L \rrbracket$
 $\implies P \ (x \sqcup y)$
 $\langle \text{proof} \rangle$

lemma (**in** *lattice*) *join-closed* [*simp*]:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqcup y \in \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma (**in** *partial-order*) *sup-of-singletonI*:
 $x \in \text{carrier } L \implies \text{least } L \ x \ (\text{Upper } L \ \{x\})$
 $\langle \text{proof} \rangle$

lemma (**in** *partial-order*) *sup-of-singleton* [*simp*]:
 $x \in \text{carrier } L \implies \bigsqcup \{x\} = x$
 $\langle \text{proof} \rangle$

Condition on A : supremum exists.

lemma (**in** *lattice*) *sup-insertI*:
 $\llbracket \exists s. \text{least } L \ s \ (\text{Upper } L \ (\text{insert } x \ A)) \rrbracket \implies P \ s;$
 $\text{least } L \ a \ (\text{Upper } L \ A); x \in \text{carrier } L; A \subseteq \text{carrier } L \rrbracket$
 $\implies P \ (\bigsqcup (\text{insert } x \ A))$
 $\langle \text{proof} \rangle$

lemma (**in** *lattice*) *finite-sup-least*:

$\llbracket \text{finite } A; A \subseteq \text{carrier } L; A \sim = \{\} \rrbracket \implies \text{least } L (\bigsqcup A) (\text{Upper } L A)$
 $\langle \text{proof} \rangle$

lemma (in lattice) *finite-sup-insertI*:
assumes $P: \llbracket l. \text{least } L l (\text{Upper } L (\text{insert } x A)) \rrbracket \implies P l$
and $xA: \text{finite } A \quad x \in \text{carrier } L \quad A \subseteq \text{carrier } L$
shows $P (\bigsqcup (\text{insert } x A))$
 $\langle \text{proof} \rangle$

lemma (in lattice) *finite-sup-closed*:
 $\llbracket \text{finite } A; A \subseteq \text{carrier } L; A \sim = \{\} \rrbracket \implies \bigsqcup A \in \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma (in lattice) *join-left*:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqsubseteq x \sqcup y$
 $\langle \text{proof} \rangle$

lemma (in lattice) *join-right*:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies y \sqsubseteq x \sqcup y$
 $\langle \text{proof} \rangle$

lemma (in lattice) *sup-of-two-least*:
 $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies \text{least } L (\bigsqcup \{x, y\}) (\text{Upper } L \{x, y\})$
 $\langle \text{proof} \rangle$

lemma (in lattice) *join-le*:
assumes $\text{sub}: x \sqsubseteq z \quad y \sqsubseteq z$
and $x: x \in \text{carrier } L$ **and** $y: y \in \text{carrier } L$ **and** $z: z \in \text{carrier } L$
shows $x \sqcup y \sqsubseteq z$
 $\langle \text{proof} \rangle$

lemma (in lattice) *join-assoc-lemma*:
assumes $L: x \in \text{carrier } L \quad y \in \text{carrier } L \quad z \in \text{carrier } L$
shows $x \sqcup (y \sqcup z) = \bigsqcup \{x, y, z\}$
 $\langle \text{proof} \rangle$

lemma *join-comm*:
fixes L (**structure**)
shows $x \sqcup y = y \sqcup x$
 $\langle \text{proof} \rangle$

lemma (in lattice) *join-assoc*:
assumes $L: x \in \text{carrier } L \quad y \in \text{carrier } L \quad z \in \text{carrier } L$
shows $(x \sqcup y) \sqcup z = x \sqcup (y \sqcup z)$
 $\langle \text{proof} \rangle$

1.2.2 Infimum

lemma (in lattice) *meetI*:

$$\begin{aligned} & \llbracket \text{!!}i. \text{greatest } L \ i \ (\text{Lower } L \ \{x, y\}) \implies P \ i; \\ & x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \\ & \implies P \ (x \sqcap y) \\ & \langle \text{proof} \rangle \end{aligned}$$

lemma (in lattice) *meet-closed* [simp]:

$$\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqcap y \in \text{carrier } L$$

$$\langle \text{proof} \rangle$$

lemma (in partial-order) *inf-of-singletonI*:

$$x \in \text{carrier } L \implies \text{greatest } L \ x \ (\text{Lower } L \ \{x\})$$

$$\langle \text{proof} \rangle$$

lemma (in partial-order) *inf-of-singleton* [simp]:

$$x \in \text{carrier } L \implies \sqcap \ \{x\} = x$$

$$\langle \text{proof} \rangle$$

Condition on A: infimum exists.

lemma (in lattice) *inf-insertI*:

$$\begin{aligned} & \llbracket \text{!!}i. \text{greatest } L \ i \ (\text{Lower } L \ (\text{insert } x \ A)) \implies P \ i; \\ & \text{greatest } L \ a \ (\text{Lower } L \ A); x \in \text{carrier } L; A \subseteq \text{carrier } L \rrbracket \\ & \implies P \ (\sqcap (\text{insert } x \ A)) \\ & \langle \text{proof} \rangle \end{aligned}$$

lemma (in lattice) *finite-inf-greatest*:

$$\llbracket \text{finite } A; A \subseteq \text{carrier } L; A \sim = \{\} \rrbracket \implies \text{greatest } L \ (\sqcap A) \ (\text{Lower } L \ A)$$

$$\langle \text{proof} \rangle$$

lemma (in lattice) *finite-inf-insertI*:
assumes $P: \text{!!}i. \text{greatest } L \ i \ (\text{Lower } L \ (\text{insert } x \ A)) \implies P \ i$
and $xA: \text{finite } A \ x \in \text{carrier } L \ A \subseteq \text{carrier } L$
shows $P \ (\sqcap (\text{insert } x \ A))$

$$\langle \text{proof} \rangle$$

lemma (in lattice) *finite-inf-closed*:

$$\llbracket \text{finite } A; A \subseteq \text{carrier } L; A \sim = \{\} \rrbracket \implies \sqcap A \in \text{carrier } L$$

$$\langle \text{proof} \rangle$$

lemma (in lattice) *meet-left*:

$$\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqcap y \sqsubseteq x$$

$$\langle \text{proof} \rangle$$

lemma (in lattice) *meet-right*:

$$\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqcap y \sqsubseteq y$$

$$\langle \text{proof} \rangle$$

lemma (in lattice) *inf-of-two-greatest*:

$$\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies$$

$$\text{greatest } L \ (\sqcap \ \{x, y\}) \ (\text{Lower } L \ \{x, y\})$$

<proof>

lemma (in *lattice*) *meet-le*:

assumes *sub*: $z \sqsubseteq x$ $z \sqsubseteq y$

and $x \in \text{carrier } L$ **and** $y \in \text{carrier } L$ **and** $z \in \text{carrier } L$

shows $z \sqsubseteq x \sqcap y$

<proof>

lemma (in *lattice*) *meet-assoc-lemma*:

assumes L : $x \in \text{carrier } L$ $y \in \text{carrier } L$ $z \in \text{carrier } L$

shows $x \sqcap (y \sqcap z) = \sqcap \{x, y, z\}$

<proof>

lemma *meet-comm*:

fixes L (**structure**)

shows $x \sqcap y = y \sqcap x$

<proof>

lemma (in *lattice*) *meet-assoc*:

assumes L : $x \in \text{carrier } L$ $y \in \text{carrier } L$ $z \in \text{carrier } L$

shows $(x \sqcap y) \sqcap z = x \sqcap (y \sqcap z)$

<proof>

1.3 Total Orders

locale *total-order* = *partial-order* +

assumes *total*: $\llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqsubseteq y \mid y \sqsubseteq x$

Introduction rule: the usual definition of total order

lemma (in *partial-order*) *total-orderI*:

assumes *total*: $\forall x y. \llbracket x \in \text{carrier } L; y \in \text{carrier } L \rrbracket \implies x \sqsubseteq y \mid y \sqsubseteq x$

shows *total-order* L

<proof>

Total orders are lattices.

interpretation *total-order* < *lattice*

<proof>

1.4 Complete lattices

locale *complete-lattice* = *lattice* +

assumes *sup-exists*:

$\llbracket A \subseteq \text{carrier } L \rrbracket \implies \text{EX } s. \text{ least } L s \text{ (Upper } L A)$

and *inf-exists*:

$\llbracket A \subseteq \text{carrier } L \rrbracket \implies \text{EX } i. \text{ greatest } L i \text{ (Lower } L A)$

Introduction rule: the usual definition of complete lattice

lemma (in *partial-order*) *complete-latticeI*:

assumes *sup-exists*:

!!A. $[A \subseteq \text{carrier } L] \implies \text{EX } s. \text{ least } L \ s \ (\text{Upper } L \ A)$
and *inf-exists*:
!!A. $[A \subseteq \text{carrier } L] \implies \text{EX } i. \text{ greatest } L \ i \ (\text{Lower } L \ A)$
shows *complete-lattice* L
 $\langle \text{proof} \rangle$

constdefs (**structure** L)
 $\text{top} :: - \implies 'a \ (\top_1)$
 $\top == \text{sup } L \ (\text{carrier } L)$

 $\text{bottom} :: - \implies 'a \ (\perp_1)$
 $\perp == \text{inf } L \ (\text{carrier } L)$

lemma (**in** *complete-lattice*) *supI*:
 $[!!l. \text{ least } L \ l \ (\text{Upper } L \ A) \implies P \ l; A \subseteq \text{carrier } L]$
 $\implies P \ (\bigsqcup A)$
 $\langle \text{proof} \rangle$

lemma (**in** *complete-lattice*) *sup-closed* [*simp*]:
 $A \subseteq \text{carrier } L \implies \bigsqcup A \in \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma (**in** *complete-lattice*) *top-closed* [*simp*, *intro*]:
 $\top \in \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma (**in** *complete-lattice*) *infI*:
 $[!!i. \text{ greatest } L \ i \ (\text{Lower } L \ A) \implies P \ i; A \subseteq \text{carrier } L]$
 $\implies P \ (\bigsqcap A)$
 $\langle \text{proof} \rangle$

lemma (**in** *complete-lattice*) *inf-closed* [*simp*]:
 $A \subseteq \text{carrier } L \implies \bigsqcap A \in \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma (**in** *complete-lattice*) *bottom-closed* [*simp*, *intro*]:
 $\perp \in \text{carrier } L$
 $\langle \text{proof} \rangle$

Jacobson: Theorem 8.1

lemma *Lower-empty* [*simp*]:
 $\text{Lower } L \ \{\} = \text{carrier } L$
 $\langle \text{proof} \rangle$

lemma *Upper-empty* [*simp*]:
 $\text{Upper } L \ \{\} = \text{carrier } L$
 $\langle \text{proof} \rangle$

theorem (in *partial-order*) *complete-lattice-criterion1*:
assumes *top-exists*: *EX g. greatest L g (carrier L)*
and *inf-exists*:
 !!A. [| *A* \subseteq *carrier L*; *A* \sim = {} |] ==> *EX i. greatest L i (Lower L A)*
shows *complete-lattice L*
 <proof>

1.5 Examples

1.5.1 Powerset of a Set is a Complete Lattice

theorem *powerset-is-complete-lattice*:
complete-lattice (| carrier = Pow A, le = op \subseteq |)
 (is *complete-lattice ?L*)
 <proof>

An other example, that of the lattice of subgroups of a group, can be found in Group theory (Section 2.7).

end

theory *Group imports FuncSet Lattice begin*

2 Monoids and Groups

2.1 Definitions

Definitions follow [2].

record *'a monoid* = *'a partial-object* +
mult :: [*'a*, *'a*] \Rightarrow *'a* (**infixl** \otimes_1 70)
one :: *'a* (**1**)

constdefs (structure *G*)
m-inv :: (*'a*, *'b*) *monoid-scheme* \Rightarrow *'a* \Rightarrow *'a* (*inv1* - [81] 80)
inv x == (*THE y. y* \in *carrier G* & *x* \otimes *y* = **1** & *y* \otimes *x* = **1**)

Units :: - \Rightarrow *'a set*

— The set of invertible elements

Units G == {*y. y* \in *carrier G* & ($\exists x \in$ *carrier G. x* \otimes *y* = **1** & *y* \otimes *x* = **1**)}

consts

pow :: [(*'a*, *'m*) *monoid-scheme*, *'a*, *'b::number*] \Rightarrow *'a* (**infixr** $'(^)1$ 75)

defs (overloaded)

nat-pow-def: *pow G a n* == *nat-rec 1_G (%u b. b* \otimes_G *a) n*

int-pow-def: *pow G a z* ==

let p = *nat-rec 1_G (%u b. b* \otimes_G *a)*

in if neg z then inv_G (p (nat (-z))) else p (nat z)

locale monoid =
fixes G (**structure**)
assumes *m-closed* [*intro, simp*]:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies x \otimes y \in \text{carrier } G$
and *m-assoc*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket$
 $\implies (x \otimes y) \otimes z = x \otimes (y \otimes z)$
and *one-closed* [*intro, simp*]: $\mathbf{1} \in \text{carrier } G$
and *l-one* [*simp*]: $x \in \text{carrier } G \implies \mathbf{1} \otimes x = x$
and *r-one* [*simp*]: $x \in \text{carrier } G \implies x \otimes \mathbf{1} = x$

lemma monoidI:
fixes G (**structure**)
assumes *m-closed*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies x \otimes y \in \text{carrier } G$
and *one-closed*: $\mathbf{1} \in \text{carrier } G$
and *m-assoc*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$
 $(x \otimes y) \otimes z = x \otimes (y \otimes z)$
and *l-one*: $\llbracket x \in \text{carrier } G \rrbracket \implies \mathbf{1} \otimes x = x$
and *r-one*: $\llbracket x \in \text{carrier } G \rrbracket \implies x \otimes \mathbf{1} = x$
shows *monoid* G
 $\langle \text{proof} \rangle$

lemma (in monoid) Units-closed [*dest*]:
 $x \in \text{Units } G \implies x \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (in monoid) inv-unique:
assumes *eq*: $y \otimes x = \mathbf{1} \quad x \otimes y' = \mathbf{1}$
and G : $x \in \text{carrier } G \quad y \in \text{carrier } G \quad y' \in \text{carrier } G$
shows $y = y'$
 $\langle \text{proof} \rangle$

lemma (in monoid) Units-one-closed [*intro, simp*]:
 $\mathbf{1} \in \text{Units } G$
 $\langle \text{proof} \rangle$

lemma (in monoid) Units-inv-closed [*intro, simp*]:
 $x \in \text{Units } G \implies \text{inv } x \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (in monoid) Units-l-inv-ex:
 $x \in \text{Units } G \implies \exists y \in \text{carrier } G. y \otimes x = \mathbf{1}$
 $\langle \text{proof} \rangle$

lemma (in monoid) Units-r-inv-ex:

$x \in \text{Units } G \implies \exists y \in \text{carrier } G. x \otimes y = \mathbf{1}$
 ⟨proof⟩

lemma (in monoid) *Units-l-inv*:
 $x \in \text{Units } G \implies \text{inv } x \otimes x = \mathbf{1}$
 ⟨proof⟩

lemma (in monoid) *Units-r-inv*:
 $x \in \text{Units } G \implies x \otimes \text{inv } x = \mathbf{1}$
 ⟨proof⟩

lemma (in monoid) *Units-inv-Units* [intro, simp]:
 $x \in \text{Units } G \implies \text{inv } x \in \text{Units } G$
 ⟨proof⟩

lemma (in monoid) *Units-l-cancel* [simp]:
 [[$x \in \text{Units } G; y \in \text{carrier } G; z \in \text{carrier } G$]] \implies
 $(x \otimes y = x \otimes z) = (y = z)$
 ⟨proof⟩

lemma (in monoid) *Units-inv-inv* [simp]:
 $x \in \text{Units } G \implies \text{inv } (\text{inv } x) = x$
 ⟨proof⟩

lemma (in monoid) *inv-inj-on-Units*:
 $\text{inj-on } (m\text{-inv } G) (\text{Units } G)$
 ⟨proof⟩

lemma (in monoid) *Units-inv-comm*:
assumes $\text{inv}: x \otimes y = \mathbf{1}$
and $G: x \in \text{Units } G \ y \in \text{Units } G$
shows $y \otimes x = \mathbf{1}$
 ⟨proof⟩

Power

lemma (in monoid) *nat-pow-closed* [intro, simp]:
 $x \in \text{carrier } G \implies x (^) (n::\text{nat}) \in \text{carrier } G$
 ⟨proof⟩

lemma (in monoid) *nat-pow-0* [simp]:
 $x (^) (0::\text{nat}) = \mathbf{1}$
 ⟨proof⟩

lemma (in monoid) *nat-pow-Suc* [simp]:
 $x (^) (\text{Suc } n) = x (^) n \otimes x$
 ⟨proof⟩

lemma (in monoid) *nat-pow-one* [simp]:
 $\mathbf{1} (^) (n::\text{nat}) = \mathbf{1}$

<proof>

lemma (in monoid) *nat-pow-mult*:

$x \in \text{carrier } G \implies x (^) (n::\text{nat}) \otimes x (^) m = x (^) (n + m)$

<proof>

lemma (in monoid) *nat-pow-pow*:

$x \in \text{carrier } G \implies (x (^) n) (^) m = x (^) (n * m::\text{nat})$

<proof>

A group is a monoid all of whose elements are invertible.

locale *group = monoid +*

assumes *Units: carrier G <= Units G*

lemma (in group) *is-group: group G <proof>*

theorem *groupI*:

fixes *G (structure)*

assumes *m-closed [simp]*:

$\forall x y. [| x \in \text{carrier } G; y \in \text{carrier } G |] \implies x \otimes y \in \text{carrier } G$

and *one-closed [simp]: 1 ∈ carrier G*

and *m-assoc*:

$\forall x y z. [| x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G |] \implies$

$(x \otimes y) \otimes z = x \otimes (y \otimes z)$

and *l-one [simp]: !x. x ∈ carrier G ==> 1 ⊗ x = x*

and *l-inv-ex: !x. x ∈ carrier G ==> ∃ y ∈ carrier G. y ⊗ x = 1*

shows *group G*

<proof>

lemma (in monoid) *monoid-groupI*:

assumes *l-inv-ex*:

$\forall x. x \in \text{carrier } G \implies \exists y \in \text{carrier } G. y \otimes x = \mathbf{1}$

shows *group G*

<proof>

lemma (in group) *Units-eq [simp]*:

$\text{Units } G = \text{carrier } G$

<proof>

lemma (in group) *inv-closed [intro, simp]*:

$x \in \text{carrier } G \implies \text{inv } x \in \text{carrier } G$

<proof>

lemma (in group) *l-inv-ex [simp]*:

$x \in \text{carrier } G \implies \exists y \in \text{carrier } G. y \otimes x = \mathbf{1}$

<proof>

lemma (in group) *r-inv-ex [simp]*:

$x \in \text{carrier } G \implies \exists y \in \text{carrier } G. x \otimes y = \mathbf{1}$
 ⟨proof⟩

lemma (in group) *l-inv* [simp]:
 $x \in \text{carrier } G \implies \text{inv } x \otimes x = \mathbf{1}$
 ⟨proof⟩

2.2 Cancellation Laws and Basic Properties

lemma (in group) *l-cancel* [simp]:
 $[[x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G]] \implies$
 $(x \otimes y = x \otimes z) = (y = z)$
 ⟨proof⟩

lemma (in group) *r-inv* [simp]:
 $x \in \text{carrier } G \implies x \otimes \text{inv } x = \mathbf{1}$
 ⟨proof⟩

lemma (in group) *r-cancel* [simp]:
 $[[x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G]] \implies$
 $(y \otimes x = z \otimes x) = (y = z)$
 ⟨proof⟩

lemma (in group) *inv-one* [simp]:
 $\text{inv } \mathbf{1} = \mathbf{1}$
 ⟨proof⟩

lemma (in group) *inv-inv* [simp]:
 $x \in \text{carrier } G \implies \text{inv } (\text{inv } x) = x$
 ⟨proof⟩

lemma (in group) *inv-inj*:
 $\text{inj-on } (m\text{-inv } G) (\text{carrier } G)$
 ⟨proof⟩

lemma (in group) *inv-mult-group*:
 $[[x \in \text{carrier } G; y \in \text{carrier } G]] \implies \text{inv } (x \otimes y) = \text{inv } y \otimes \text{inv } x$
 ⟨proof⟩

lemma (in group) *inv-comm*:
 $[[x \otimes y = \mathbf{1}; x \in \text{carrier } G; y \in \text{carrier } G]] \implies y \otimes x = \mathbf{1}$
 ⟨proof⟩

lemma (in group) *inv-equality*:
 $[[y \otimes x = \mathbf{1}; x \in \text{carrier } G; y \in \text{carrier } G]] \implies \text{inv } x = y$
 ⟨proof⟩

Power

lemma (in group) *int-pow-def2*:

$a (^) (z::int) = (if\ neg\ z\ then\ inv\ (a\ (^)\ (nat\ (-z)))\ else\ a\ (^)\ (nat\ z))$
 $\langle proof \rangle$

lemma (in group) *int-pow-0* [simp]:
 $x (^) (0::int) = \mathbf{1}$
 $\langle proof \rangle$

lemma (in group) *int-pow-one* [simp]:
 $\mathbf{1} (^) (z::int) = \mathbf{1}$
 $\langle proof \rangle$

2.3 Subgroups

locale *subgroup* =
fixes H and G (**structure**)
assumes *subset*: $H \subseteq carrier\ G$
and *m-closed* [intro, simp]: $\llbracket x \in H; y \in H \rrbracket \implies x \otimes y \in H$
and *one-closed* [simp]: $\mathbf{1} \in H$
and *m-inv-closed* [intro,simp]: $x \in H \implies inv\ x \in H$

lemma (in subgroup) *is-subgroup*:
subgroup $H\ G$ $\langle proof \rangle$

declare (in subgroup) *group.intro* [intro]

lemma (in subgroup) *mem-carrier* [simp]:
 $x \in H \implies x \in carrier\ G$
 $\langle proof \rangle$

lemma *subgroup-imp-subset*:
subgroup $H\ G \implies H \subseteq carrier\ G$
 $\langle proof \rangle$

lemma (in subgroup) *subgroup-is-group* [intro]:
includes *group* G
shows *group* ($G(\backslash carrier := H)$)
 $\langle proof \rangle$

Since H is nonempty, it contains some element x . Since it is closed under inverse, it contains $inv\ x$. Since it is closed under product, it contains $x \otimes inv\ x = \mathbf{1}$.

lemma (in group) *one-in-subset*:
 $\llbracket H \subseteq carrier\ G; H \neq \{\}; \forall a \in H. inv\ a \in H; \forall a \in H. \forall b \in H. a \otimes b \in H \rrbracket$
 $\implies \mathbf{1} \in H$
 $\langle proof \rangle$

A characterization of subgroups: closed, non-empty subset.

lemma (in group) *subgroupI*:
assumes *subset*: $H \subseteq carrier\ G$ **and** *non-empty*: $H \neq \{\}$

```

and inv: !!a. a ∈ H ⇒ inv a ∈ H
and mult: !!a b. [a ∈ H; b ∈ H] ⇒ a ⊗ b ∈ H
shows subgroup H G
⟨proof⟩

```

```

declare monoid.one-closed [iff] group.inv-closed [simp]
monoid.l-one [simp] monoid.r-one [simp] group.inv-inv [simp]

```

```

lemma subgroup-nonempty:
~ subgroup {} G
⟨proof⟩

```

```

lemma (in subgroup) finite-imp-card-positive:
finite (carrier G) ⇒ 0 < card H
⟨proof⟩

```

2.4 Direct Products

```

constdefs
DirProd :: - ⇒ - ⇒ ('a × 'b) monoid (infixr ×× 80)
G ×× H ≡ (|carrier = carrier G × carrier H,
           mult = (λ(g, h) (g', h'). (g ⊗G g', h ⊗H h')),
           one = (1G, 1H)|)

```

```

lemma DirProd-monoid:
includes monoid G + monoid H
shows monoid (G ×× H)
⟨proof⟩

```

Does not use the previous result because it's easier just to use auto.

```

lemma DirProd-group:
includes group G + group H
shows group (G ×× H)
⟨proof⟩

```

```

lemma carrier-DirProd [simp]:
carrier (G ×× H) = carrier G × carrier H
⟨proof⟩

```

```

lemma one-DirProd [simp]:
1G ×× H = (1G, 1H)
⟨proof⟩

```

```

lemma mult-DirProd [simp]:
(g, h) ⊗(G ×× H) (g', h') = (g ⊗G g', h ⊗H h')
⟨proof⟩

```

```

lemma inv-DirProd [simp]:
includes group G + group H

```

```

assumes  $g: g \in \text{carrier } G$ 
and  $h: h \in \text{carrier } H$ 
shows  $m\text{-inv } (G \times\times H) (g, h) = (\text{inv}_G g, \text{inv}_H h)$ 
<proof>

```

This alternative proof of the previous result demonstrates `interpret`. It uses `Prod.inv-equality` (available after `interpret`) instead of `group.inv-equality` [`OF DirProd-group`].

```

lemma
includes  $\text{group } G + \text{group } H$ 
assumes  $g: g \in \text{carrier } G$ 
and  $h: h \in \text{carrier } H$ 
shows  $m\text{-inv } (G \times\times H) (g, h) = (\text{inv}_G g, \text{inv}_H h)$ 
<proof>

```

2.5 Homomorphisms and Isomorphisms

```

constdefs (structure  $G$  and  $H$ )
   $\text{hom} :: - \Rightarrow - \Rightarrow ('a \Rightarrow 'b) \text{ set}$ 
   $\text{hom } G H ==$ 
     $\{h. h \in \text{carrier } G \rightarrow \text{carrier } H \ \&$ 
       $(\forall x \in \text{carrier } G. \forall y \in \text{carrier } G. h (x \otimes_G y) = h x \otimes_H h y)\}$ 

```

```

lemma hom-mult:
   $[[ h \in \text{hom } G H; x \in \text{carrier } G; y \in \text{carrier } G ]]$ 
   $==> h (x \otimes_G y) = h x \otimes_H h y$ 
<proof>

```

```

lemma hom-closed:
   $[[ h \in \text{hom } G H; x \in \text{carrier } G ]]$   $==> h x \in \text{carrier } H$ 
<proof>

```

```

lemma (in group) hom-compose:
   $[[ h \in \text{hom } G H; i \in \text{hom } H I ]]$   $==> \text{compose } (\text{carrier } G) i h \in \text{hom } G I$ 
<proof>

```

```

constdefs
   $\text{iso} :: - \Rightarrow - \Rightarrow ('a \Rightarrow 'b) \text{ set}$  (infixr  $\cong$  60)
   $G \cong H == \{h. h \in \text{hom } G H \ \& \ \text{bij-betw } h (\text{carrier } G) (\text{carrier } H)\}$ 

```

```

lemma iso-refl:  $(\%x. x) \in G \cong G$ 
<proof>

```

```

lemma (in group) iso-sym:
   $h \in G \cong H \implies \text{Inv } (\text{carrier } G) h \in H \cong G$ 
<proof>

```

```

lemma (in group) iso-trans:
   $[[ h \in G \cong H; i \in H \cong I ]]$   $==> (\text{compose } (\text{carrier } G) i h) \in G \cong I$ 

```

$\langle proof \rangle$

lemma *DirProd-commute-iso*:

shows $(\lambda(x,y). (y,x)) \in (G \times \times H) \cong (H \times \times G)$

$\langle proof \rangle$

lemma *DirProd-assoc-iso*:

shows $(\lambda(x,y,z). (x,(y,z))) \in (G \times \times H \times \times I) \cong (G \times \times (H \times \times I))$

$\langle proof \rangle$

Basis for homomorphism proofs: we assume two groups G and H , with a homomorphism h between them

locale *group-hom* = *group* G + *group* H + *var* h +

assumes *homh*: $h \in \text{hom } G \ H$

notes *hom-mult* [*simp*] = *hom-mult* [*OF homh*]

and *hom-closed* [*simp*] = *hom-closed* [*OF homh*]

lemma (**in** *group-hom*) *one-closed* [*simp*]:

$h \ \mathbf{1} \in \text{carrier } H$

$\langle proof \rangle$

lemma (**in** *group-hom*) *hom-one* [*simp*]:

$h \ \mathbf{1} = \mathbf{1}_H$

$\langle proof \rangle$

lemma (**in** *group-hom*) *inv-closed* [*simp*]:

$x \in \text{carrier } G \implies h \ (\text{inv } x) \in \text{carrier } H$

$\langle proof \rangle$

lemma (**in** *group-hom*) *hom-inv* [*simp*]:

$x \in \text{carrier } G \implies h \ (\text{inv } x) = \text{inv}_H (h \ x)$

$\langle proof \rangle$

2.6 Commutative Structures

Naming convention: multiplicative structures that are commutative are called *commutative*, additive structures are called *Abelian*.

locale *comm-monoid* = *monoid* +

assumes *m-comm*: $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies x \otimes y = y \otimes x$

lemma (**in** *comm-monoid*) *m-lcomm*:

$\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$

$x \otimes (y \otimes z) = y \otimes (x \otimes z)$

$\langle proof \rangle$

lemmas (**in** *comm-monoid*) *m-ac* = *m-assoc* *m-comm* *m-lcomm*

lemma *comm-monoidI*:

fixes G (**structure**)

assumes m -closed:

!! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==> $x \otimes y \in \text{carrier } G$

and one -closed: $\mathbf{1} \in \text{carrier } G$

and m -assoc:

!! $x y z$. [$x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G$] ==>

$(x \otimes y) \otimes z = x \otimes (y \otimes z)$

and l -one: !! x . $x \in \text{carrier } G$ ==> $\mathbf{1} \otimes x = x$

and m -comm:

!! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==> $x \otimes y = y \otimes x$

shows $comm$ -monoid G

$\langle proof \rangle$

lemma (**in** $monoid$) $monoid$ - $comm$ - $monoidI$:

assumes m -comm:

!! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==> $x \otimes y = y \otimes x$

shows $comm$ -monoid G

$\langle proof \rangle$

lemma (**in** $comm$ -monoid) nat - pow - $distr$:

[$x \in \text{carrier } G; y \in \text{carrier } G$] ==>

$(x \otimes y) (^) (n::nat) = x (^) n \otimes y (^) n$

$\langle proof \rangle$

locale $comm$ -group = $comm$ -monoid + group

lemma (**in** group) $group$ - $comm$ - $groupI$:

assumes m -comm: !! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==>

$x \otimes y = y \otimes x$

shows $comm$ -group G

$\langle proof \rangle$

lemma $comm$ - $groupI$:

fixes G (**structure**)

assumes m -closed:

!! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==> $x \otimes y \in \text{carrier } G$

and one -closed: $\mathbf{1} \in \text{carrier } G$

and m -assoc:

!! $x y z$. [$x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G$] ==>

$(x \otimes y) \otimes z = x \otimes (y \otimes z)$

and m -comm:

!! $x y$. [$x \in \text{carrier } G; y \in \text{carrier } G$] ==> $x \otimes y = y \otimes x$

and l -one: !! x . $x \in \text{carrier } G$ ==> $\mathbf{1} \otimes x = x$

and l -inv- ex : !! x . $x \in \text{carrier } G$ ==> $\exists y \in \text{carrier } G. y \otimes x = \mathbf{1}$

shows $comm$ -group G

$\langle proof \rangle$

lemma (in *comm-group*) *inv-mult*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies \text{inv } (x \otimes y) = \text{inv } x \otimes \text{inv } y$
 ⟨proof⟩

2.7 The Lattice of Subgroups of a Group

theorem (in *group*) *subgroups-partial-order*:
 $\text{partial-order } (\llbracket \text{carrier} = \{H. \text{subgroup } H \ G\}, \text{le} = \text{op } \subseteq \rrbracket)$
 ⟨proof⟩

lemma (in *group*) *subgroup-self*:
 $\text{subgroup } (\text{carrier } G) \ G$
 ⟨proof⟩

lemma (in *group*) *subgroup-imp-group*:
 $\text{subgroup } H \ G \implies \text{group } (G (\llbracket \text{carrier} := H \rrbracket))$
 ⟨proof⟩

lemma (in *group*) *is-monoid [intro, simp]*:
 $\text{monoid } G$
 ⟨proof⟩

lemma (in *group*) *subgroup-inv-equality*:
 $\llbracket \text{subgroup } H \ G; x \in H \rrbracket \implies \text{m-inv } (G (\llbracket \text{carrier} := H \rrbracket)) \ x = \text{inv } x$
 ⟨proof⟩

theorem (in *group*) *subgroups-Inter*:
assumes *subgr*: $(\forall H. H \in A \implies \text{subgroup } H \ G)$
and *not-empty*: $A \neq \{\}$
shows $\text{subgroup } (\bigcap A) \ G$
 ⟨proof⟩

theorem (in *group*) *subgroups-complete-lattice*:
 $\text{complete-lattice } (\llbracket \text{carrier} = \{H. \text{subgroup } H \ G\}, \text{le} = \text{op } \subseteq \rrbracket)$
 (is complete-lattice ?L)
 ⟨proof⟩

end

theory *FiniteProduct* **imports** *Group* **begin**

3 Product Operator for Commutative Monoids

3.1 Inductive Definition of a Relation for Products over Sets

Instantiation of locale *LC* of theory *Finite-Set* is not possible, because here we have explicit typing rules like $x \in \text{carrier } G$. We introduce an explicit

argument for the domain D .

inductive-set

$foldSetD :: ['a\ set, 'b \Rightarrow 'a \Rightarrow 'a, 'a] \Rightarrow ('b\ set * 'a)\ set$
for $D :: 'a\ set$ **and** $f :: 'b \Rightarrow 'a \Rightarrow 'a$ **and** $e :: 'a$
where
 $emptyI$ [intro]: $e \in D \Rightarrow (\{\}, e) \in foldSetD\ D\ f\ e$
 $insertI$ [intro]: $[[\ x \sim: A; f\ x\ y \in D; (A, y) \in foldSetD\ D\ f\ e\] \Rightarrow$
 $(insert\ x\ A, f\ x\ y) \in foldSetD\ D\ f\ e$

inductive-cases $empty-foldSetDE$ [elim!]: $(\{\}, x) \in foldSetD\ D\ f\ e$

constdefs

$foldD :: ['a\ set, 'b \Rightarrow 'a \Rightarrow 'a, 'a, 'b\ set] \Rightarrow 'a$
 $foldD\ D\ f\ e\ A == THE\ x.\ (A, x) \in foldSetD\ D\ f\ e$

lemma $foldSetD-closed$:

$[[\ (A, z) \in foldSetD\ D\ f\ e ; e \in D; !!x\ y.\ [[\ x \in A; y \in D\] \Rightarrow f\ x\ y \in D$
 $]] \Rightarrow z \in D$
 $\langle proof \rangle$

lemma $Diff1-foldSetD$:

$[[\ (A - \{x\}, y) \in foldSetD\ D\ f\ e; x \in A; f\ x\ y \in D\] \Rightarrow$
 $(A, f\ x\ y) \in foldSetD\ D\ f\ e$
 $\langle proof \rangle$

lemma $foldSetD-imp-finite$ [simp]: $(A, x) \in foldSetD\ D\ f\ e \Rightarrow finite\ A$

$\langle proof \rangle$

lemma $finite-imp-foldSetD$:

$[[\ finite\ A; e \in D; !!x\ y.\ [[\ x \in A; y \in D\] \Rightarrow f\ x\ y \in D\] \Rightarrow$
 $EX\ x.\ (A, x) \in foldSetD\ D\ f\ e$
 $\langle proof \rangle$

3.2 Left-Commutative Operations

locale $LCD =$

fixes $B :: 'b\ set$
and $D :: 'a\ set$
and $f :: 'b \Rightarrow 'a \Rightarrow 'a$ (**infixl** \cdot 70)
assumes $left-commute$:
 $[[\ x \in B; y \in B; z \in D\] \Rightarrow x \cdot (y \cdot z) = y \cdot (x \cdot z)$
and $f-closed$ [simp, intro!]: $!!x\ y.\ [[\ x \in B; y \in D\] \Rightarrow f\ x\ y \in D$

lemma (**in** LCD) $foldSetD-closed$ [dest]:

$(A, z) \in foldSetD\ D\ f\ e \Rightarrow z \in D$
 $\langle proof \rangle$

lemma (**in** LCD) $Diff1-foldSetD$:

$[[\ (A - \{x\}, y) \in foldSetD\ D\ f\ e; x \in A; A \subseteq B\] \Rightarrow$

$(A, f x y) \in \text{foldSetD } D f e$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldSetD-imp-finite* [*simp*]:
 $(A, x) \in \text{foldSetD } D f e \implies \text{finite } A$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *finite-imp-foldSetD*:
 $\llbracket \text{finite } A; A \subseteq B; e \in D \rrbracket \implies \exists x. (A, x) \in \text{foldSetD } D f e$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldSetD-determ-aux*:
 $e \in D \implies \forall A x. A \subseteq B \ \& \ \text{card } A < n \implies (A, x) \in \text{foldSetD } D f e \implies$
 $(\forall y. (A, y) \in \text{foldSetD } D f e \implies y = x)$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldSetD-determ*:
 $\llbracket (A, x) \in \text{foldSetD } D f e; (A, y) \in \text{foldSetD } D f e; e \in D; A \subseteq B \rrbracket$
 $\implies y = x$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldD-equality*:
 $\llbracket (A, y) \in \text{foldSetD } D f e; e \in D; A \subseteq B \rrbracket \implies \text{foldD } D f e A = y$
 $\langle \text{proof} \rangle$

lemma *foldD-empty* [*simp*]:
 $e \in D \implies \text{foldD } D f e \{\} = e$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldD-insert-aux*:
 $\llbracket x \sim: A; x \in B; e \in D; A \subseteq B \rrbracket \implies$
 $((\text{insert } x A, v) \in \text{foldSetD } D f e) =$
 $(\exists y. (A, y) \in \text{foldSetD } D f e \ \& \ v = f x y)$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldD-insert*:
 $\llbracket \text{finite } A; x \sim: A; x \in B; e \in D; A \subseteq B \rrbracket \implies$
 $\text{foldD } D f e (\text{insert } x A) = f x (\text{foldD } D f e A)$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldD-closed* [*simp*]:
 $\llbracket \text{finite } A; e \in D; A \subseteq B \rrbracket \implies \text{foldD } D f e A \in D$
 $\langle \text{proof} \rangle$

lemma (in *LCD*) *foldD-commute*:
 $\llbracket \text{finite } A; x \in B; e \in D; A \subseteq B \rrbracket \implies$
 $f x (\text{foldD } D f e A) = \text{foldD } D f (f x e) A$
 $\langle \text{proof} \rangle$

lemma *Int-mono2*:

$\llbracket A \subseteq C; B \subseteq C \rrbracket \implies A \text{ Int } B \subseteq C$
 $\langle \text{proof} \rangle$

lemma (**in** *LCD*) *foldD-nest-Un-Int*:

$\llbracket \text{finite } A; \text{finite } C; e \in D; A \subseteq B; C \subseteq B \rrbracket \implies$
 $\text{foldD } D f (\text{foldD } D f e C) A = \text{foldD } D f (\text{foldD } D f e (A \text{ Int } C)) (A \text{ Un } C)$
 $\langle \text{proof} \rangle$

lemma (**in** *LCD*) *foldD-nest-Un-disjoint*:

$\llbracket \text{finite } A; \text{finite } B; A \text{ Int } B = \{\}; e \in D; A \subseteq B; C \subseteq B \rrbracket$
 $\implies \text{foldD } D f e (A \text{ Un } B) = \text{foldD } D f (\text{foldD } D f e B) A$
 $\langle \text{proof} \rangle$

declare *foldSetD-imp-finite* [*simp del*]

empty-foldSetDE [*rule del*]

foldSetD.intros [*rule del*]

declare (**in** *LCD*)

foldSetD-closed [*rule del*]

3.3 Commutative Monoids

We enter a more restrictive context, with $f :: 'a \Rightarrow 'a \Rightarrow 'a$ instead of $'b \Rightarrow 'a \Rightarrow 'a$.

locale *ACeD* =

fixes $D :: 'a \text{ set}$

and $f :: 'a \Rightarrow 'a \Rightarrow 'a$ (**infixl** \cdot 70)

and $e :: 'a$

assumes *ident* [*simp*]: $x \in D \implies x \cdot e = x$

and *commute*: $\llbracket x \in D; y \in D \rrbracket \implies x \cdot y = y \cdot x$

and *assoc*: $\llbracket x \in D; y \in D; z \in D \rrbracket \implies (x \cdot y) \cdot z = x \cdot (y \cdot z)$

and *e-closed* [*simp*]: $e \in D$

and *f-closed* [*simp*]: $\llbracket x \in D; y \in D \rrbracket \implies x \cdot y \in D$

lemma (**in** *ACeD*) *left-commute*:

$\llbracket x \in D; y \in D; z \in D \rrbracket \implies x \cdot (y \cdot z) = y \cdot (x \cdot z)$
 $\langle \text{proof} \rangle$

lemmas (**in** *ACeD*) *AC = assoc commute left-commute*

lemma (**in** *ACeD*) *left-ident* [*simp*]: $x \in D \implies e \cdot x = x$

$\langle \text{proof} \rangle$

lemma (**in** *ACeD*) *foldD-Un-Int*:

$\llbracket \text{finite } A; \text{finite } B; A \subseteq D; B \subseteq D \rrbracket \implies$

$\text{foldD } D f e A \cdot \text{foldD } D f e B =$

$\text{foldD } D f e (A \text{ Un } B) \cdot \text{foldD } D f e (A \text{ Int } B)$

$\langle \text{proof} \rangle$

lemma (in *ACeD*) *foldD-Un-disjoint*:

$$\llbracket \text{finite } A; \text{finite } B; A \text{ Int } B = \{\}; A \subseteq D; B \subseteq D \rrbracket \implies$$

$$\text{foldD } D \text{ f e } (A \text{ Un } B) = \text{foldD } D \text{ f e } A \cdot \text{foldD } D \text{ f e } B$$
<proof>

3.4 Products over Finite Sets

constdefs (structure *G*)

$$\text{finprod} :: [('b, 'm) \text{ monoid-scheme}, 'a \Rightarrow 'b, 'a \text{ set}] \Rightarrow 'b$$

$$\text{finprod } G \text{ f } A == \text{if finite } A$$

$$\text{then foldD (carrier } G) (\text{mult } G \text{ o f)} \mathbf{1} \ A$$

$$\text{else arbitrary}$$

syntax

$$\text{-finprod} :: \text{index} \Rightarrow \text{idt} \Rightarrow 'a \text{ set} \Rightarrow 'b \Rightarrow 'b$$

$$((\exists \otimes \text{--} \cdot \cdot) [1000, 0, 51, 10] 10)$$

syntax (*xsymbols*)

$$\text{-finprod} :: \text{index} \Rightarrow \text{idt} \Rightarrow 'a \text{ set} \Rightarrow 'b \Rightarrow 'b$$

$$((\exists \otimes \text{--} \in \cdot \cdot) [1000, 0, 51, 10] 10)$$

syntax (*HTML output*)

$$\text{-finprod} :: \text{index} \Rightarrow \text{idt} \Rightarrow 'a \text{ set} \Rightarrow 'b \Rightarrow 'b$$

$$((\exists \otimes \text{--} \in \cdot \cdot) [1000, 0, 51, 10] 10)$$

translations

$$\otimes_{i:A}. b == \text{finprod} \diamond_1 (\%i. b) \ A$$
— Beware of argument permutation!

lemma (in *comm-monoid*) *finprod-empty* [*simp*]:

$$\text{finprod } G \text{ f } \{\} = \mathbf{1}$$
<proof>

declare *funcsetI* [*intro*]
funcset-mem [*dest*]

lemma (in *comm-monoid*) *finprod-insert* [*simp*]:

$$\llbracket \text{finite } F; a \notin F; f \in F \rightarrow \text{carrier } G; f \ a \in \text{carrier } G \rrbracket \implies$$

$$\text{finprod } G \text{ f } (\text{insert } a \ F) = f \ a \otimes \text{finprod } G \text{ f } F$$
<proof>

lemma (in *comm-monoid*) *finprod-one* [*simp*]:

$$\text{finite } A \implies (\otimes_{i:A}. \mathbf{1}) = \mathbf{1}$$
<proof>

lemma (in *comm-monoid*) *finprod-closed* [*simp*]:
fixes *A*
assumes *fn*: *finite A* **and** *f*: *f* \in *A* \rightarrow *carrier G*
shows *finprod G f A* \in *carrier G*
<proof>

lemma *funcset-Int-left* [*simp*, *intro*]:

$\llbracket f \in A \rightarrow C; f \in B \rightarrow C \rrbracket \implies f \in A \text{ Int } B \rightarrow C$
 $\langle \text{proof} \rangle$

lemma *funcset-Un-left* [*iff*]:
 $(f \in A \text{ Un } B \rightarrow C) = (f \in A \rightarrow C \ \& \ f \in B \rightarrow C)$
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-Un-Int*:
 $\llbracket \text{finite } A; \text{finite } B; g \in A \rightarrow \text{carrier } G; g \in B \rightarrow \text{carrier } G \rrbracket \implies$
 $\text{finprod } G \ g \ (A \text{ Un } B) \otimes \text{finprod } G \ g \ (A \text{ Int } B) =$
 $\text{finprod } G \ g \ A \otimes \text{finprod } G \ g \ B$
 — The reversed orientation looks more natural, but LOOPS as a simprule!
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-Un-disjoint*:
 $\llbracket \text{finite } A; \text{finite } B; A \text{ Int } B = \{\};$
 $g \in A \rightarrow \text{carrier } G; g \in B \rightarrow \text{carrier } G \rrbracket$
 $\implies \text{finprod } G \ g \ (A \text{ Un } B) = \text{finprod } G \ g \ A \otimes \text{finprod } G \ g \ B$
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-multf*:
 $\llbracket \text{finite } A; f \in A \rightarrow \text{carrier } G; g \in A \rightarrow \text{carrier } G \rrbracket \implies$
 $\text{finprod } G \ (\%x. f \ x \otimes g \ x) \ A = (\text{finprod } G \ f \ A \otimes \text{finprod } G \ g \ A)$
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-cong'*:
 $\llbracket A = B; g \in B \rightarrow \text{carrier } G;$
 $!!i. i \in B \implies f \ i = g \ i \rrbracket \implies \text{finprod } G \ f \ A = \text{finprod } G \ g \ B$
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-cong*:
 $\llbracket A = B; f \in B \rightarrow \text{carrier } G = \text{True};$
 $!!i. i \in B \implies f \ i = g \ i \rrbracket \implies \text{finprod } G \ f \ A = \text{finprod } G \ g \ B$
 $\langle \text{proof} \rangle$

Usually, if this rule causes a failed congruence proof error, the reason is that the premise $g \in B \rightarrow \text{carrier } G$ cannot be shown. Adding *Pi-def* to the simpset is often useful. For this reason, *comm-monoid.finprod-cong* is not added to the simpset by default.

declare *funcsetI* [*rule del*]
funcset-mem [*rule del*]

lemma (*in comm-monoid*) *finprod-0* [*simp*]:
 $f \in \{0::\text{nat}\} \rightarrow \text{carrier } G \implies \text{finprod } G \ f \ \{..0\} = f \ 0$
 $\langle \text{proof} \rangle$

lemma (*in comm-monoid*) *finprod-Suc* [*simp*]:
 $f \in \{..\text{Suc } n\} \rightarrow \text{carrier } G \implies$

$finprod\ G\ f\ \{..Suc\ n\} = (f\ (Suc\ n) \otimes finprod\ G\ f\ \{..n\})$
 <proof>

lemma (in *comm-monoid*) *finprod-Suc2*:
 $f \in \{..Suc\ n\} \rightarrow carrier\ G \implies$
 $finprod\ G\ f\ \{..Suc\ n\} = (finprod\ G\ (\%i.\ f\ (Suc\ i))\ \{..n\} \otimes f\ 0)$
 <proof>

lemma (in *comm-monoid*) *finprod-mult [simp]*:
 $[[\ f \in \{..n\} \rightarrow carrier\ G; g \in \{..n\} \rightarrow carrier\ G\]] \implies$
 $finprod\ G\ (\%i.\ f\ i \otimes g\ i)\ \{..n::nat\} =$
 $finprod\ G\ f\ \{..n\} \otimes finprod\ G\ g\ \{..n\}$
 <proof>

end

theory *Exponent* imports *Main Primes Binomial* begin

4 The Combinatorial Argument Underlying the First Sylow Theorem

definition *exponent* :: *nat* => *nat* => *nat* **where**
exponent *p* *s* == if prime *p* then (GREATEST *r*. p^r dvd *s*) else 0

4.1 Prime Theorems

lemma *prime-imp-one-less*: prime *p* ==> *Suc* 0 < *p*
 <proof>

lemma *prime-iff*:
 (prime *p*) = (*Suc* 0 < *p* & ($\forall a\ b.\ p\ dvd\ a*b \rightarrow (p\ dvd\ a) \mid (p\ dvd\ b)$))
 <proof>

lemma *zero-less-prime-power*: prime *p* ==> 0 < p^a
 <proof>

lemma *zero-less-card-empty*: [[finite *S*; *S* ≠ {}]] ==> 0 < card(*S*)
 <proof>

lemma *prime-dvd-cases*:
 [[$p*k\ dvd\ m*n$; prime *p*]]
 ==> ($\exists x.\ k\ dvd\ x*n \ \&\ m = p*x$) | ($\exists y.\ k\ dvd\ m*y \ \&\ n = p*y$)
 <proof>

lemma *prime-power-dvd-cases* [rule-format (no-asm)]: prime p
 $\implies \forall m n. p^c \text{ dvd } m * n \implies$
 $(\forall a b. a + b = \text{Suc } c \implies p^a \text{ dvd } m \mid p^b \text{ dvd } n)$
 <proof>

lemma *div-combine*:
 $[[\text{prime } p; \sim (p^a \text{ dvd } n); p^{a+r} \text{ dvd } n * k]]$
 $\implies p^a \text{ dvd } k$
 <proof>

lemma *Suc-le-power*: $\text{Suc } 0 < p \implies \text{Suc } n \leq p^n$
 <proof>

lemma *power-dvd-bound*: $[[p^n \text{ dvd } a; \text{Suc } 0 < p; a > 0]]$ $\implies n < a$
 <proof>

4.2 Exponent Theorems

lemma *exponent-ge* [rule-format]:
 $[[p^k \text{ dvd } n; \text{prime } p; 0 < n]]$ $\implies k \leq \text{exponent } p n$
 <proof>

lemma *power-exponent-dvd*: $s > 0 \implies (p^{\text{exponent } p s}) \text{ dvd } s$
 <proof>

lemma *power-Suc-exponent-Not-dvd*:
 $[[(p * p^{\text{exponent } p s}) \text{ dvd } s; \text{prime } p]]$ $\implies s = 0$
 <proof>

lemma *exponent-power-eq* [simp]: prime $p \implies \text{exponent } p (p^a) = a$
 <proof>

lemma *exponent-equalityI*:
 $!r::\text{nat}. (p^r \text{ dvd } a) = (p^r \text{ dvd } b) \implies \text{exponent } p a = \text{exponent } p b$
 <proof>

lemma *exponent-eq-0* [simp]: $\neg \text{prime } p \implies \text{exponent } p s = 0$
 <proof>

lemma *exponent-mult-add1*: $[[a > 0; b > 0]]$
 $\implies (\text{exponent } p a) + (\text{exponent } p b) \leq \text{exponent } p (a * b)$
 <proof>

lemma *exponent-mult-add2*: $\llbracket a > 0; b > 0 \rrbracket$
 $\implies \text{exponent } p (a * b) \leq (\text{exponent } p a) + (\text{exponent } p b)$
 $\langle \text{proof} \rangle$

lemma *exponent-mult-add*: $\llbracket a > 0; b > 0 \rrbracket$
 $\implies \text{exponent } p (a * b) = (\text{exponent } p a) + (\text{exponent } p b)$
 $\langle \text{proof} \rangle$

lemma *not-divides-exponent-0*: $\sim (p \text{ dvd } n) \implies \text{exponent } p n = 0$
 $\langle \text{proof} \rangle$

lemma *exponent-1-eq-0* [simp]: $\text{exponent } p (\text{Suc } 0) = 0$
 $\langle \text{proof} \rangle$

4.3 Main Combinatorial Argument

lemma *le-extend-mult*: $\llbracket c > 0; a \leq b \rrbracket \implies a \leq b * (c::\text{nat})$
 $\langle \text{proof} \rangle$

lemma *p-fac-forw-lemma*:
 $\llbracket (m::\text{nat}) > 0; k > 0; k < p^a; (p^r) \text{ dvd } (p^a)*m - k \rrbracket \implies r \leq a$
 $\langle \text{proof} \rangle$

lemma *p-fac-forw*: $\llbracket (m::\text{nat}) > 0; k > 0; k < p^a; (p^r) \text{ dvd } (p^a)*m - k \rrbracket$
 $\implies (p^r) \text{ dvd } (p^a) - k$
 $\langle \text{proof} \rangle$

lemma *r-le-a-forw*:
 $\llbracket (k::\text{nat}) > 0; k < p^a; p > 0; (p^r) \text{ dvd } (p^a) - k \rrbracket \implies r \leq a$
 $\langle \text{proof} \rangle$

lemma *p-fac-backw*: $\llbracket m > 0; k > 0; (p::\text{nat}) \neq 0; k < p^a; (p^r) \text{ dvd } p^a - k \rrbracket$
 $\implies (p^r) \text{ dvd } (p^a)*m - k$
 $\langle \text{proof} \rangle$

lemma *exponent-p-a-m-k-equation*: $\llbracket m > 0; k > 0; (p::\text{nat}) \neq 0; k < p^a \rrbracket$
 $\implies \text{exponent } p (p^a * m - k) = \text{exponent } p (p^a - k)$
 $\langle \text{proof} \rangle$

Suc rules that we have to delete from the simpset

lemmas *bad-Sucs = binomial-Suc-Suc mult-Suc mult-Suc-right*

lemma *p-not-div-choose-lemma* [rule-format]:
 $\llbracket \forall i. \text{Suc } i < K \dashrightarrow \text{exponent } p (\text{Suc } i) = \text{exponent } p (\text{Suc}(j+i)) \rrbracket$

$\implies k < K \dashrightarrow \text{exponent } p \text{ } ((j+k) \text{ choose } k) = 0$
 ⟨proof⟩

lemma *p-not-div-choose*:

[[$k < K$; $k \leq n$;
 $\forall j. 0 < j \ \& \ j < K \dashrightarrow \text{exponent } p \text{ } (n - k + (K - j)) = \text{exponent } p \text{ } (K - j)$]]
 $\implies \text{exponent } p \text{ } (n \text{ choose } k) = 0$
 ⟨proof⟩

lemma *const-p-fac-right*:

$m > 0 \implies \text{exponent } p \text{ } ((p \hat{a} * m - \text{Suc } 0) \text{ choose } (p \hat{a} - \text{Suc } 0)) = 0$
 ⟨proof⟩

lemma *const-p-fac*:

$m > 0 \implies \text{exponent } p \text{ } (((p \hat{a}) * m) \text{ choose } p \hat{a}) = \text{exponent } p \text{ } m$
 ⟨proof⟩

end

theory *Coset* imports *Group Exponent* begin

5 Cosets and Quotient Groups

constdefs (structure *G*)

r-coset :: $[-, 'a \text{ set}, 'a] \Rightarrow 'a \text{ set}$ (**infixl** $\#>_1$ 60)
 $H \#> a \equiv \bigcup_{h \in H}. \{h \otimes a\}$

l-coset :: $[-, 'a, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (**infixl** $<\#_1$ 60)
 $a <\# H \equiv \bigcup_{h \in H}. \{a \otimes h\}$

RCOSETS :: $[-, 'a \text{ set}] \Rightarrow ('a \text{ set})\text{set}$ (*rcosets1* - [81] 80)
 $\text{rcosets } H \equiv \bigcup_{a \in \text{carrier } G}. \{H \#> a\}$

set-mult :: $[-, 'a \text{ set}, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (**infixl** $<\#>_1$ 60)
 $H <\#> K \equiv \bigcup_{h \in H}. \bigcup_{k \in K}. \{h \otimes k\}$

SET-INV :: $[-, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (*set'-inv1* - [81] 80)
 $\text{set-inv } H \equiv \bigcup_{h \in H}. \{\text{inv } h\}$

locale *normal* = *subgroup* + *group* +

assumes *coset-eq*: $(\forall x \in \text{carrier } G. H \#> x = x <\# H)$

abbreviation

$normal\text{-}rel :: ['a\ set, ('a, 'b)\ monoid\text{-}scheme] \Rightarrow bool$ (**infixl** $\triangleleft 60$) **where**
 $H \triangleleft G \equiv normal\ H\ G$

5.1 Basic Properties of Cosets

lemma (**in group**) *coset-mult-assoc*:

$\llbracket M \subseteq carrier\ G; g \in carrier\ G; h \in carrier\ G \rrbracket$
 $\implies (M \#> g) \#> h = M \#> (g \otimes h)$

$\langle proof \rangle$

lemma (**in group**) *coset-mult-one* [*simp*]: $M \subseteq carrier\ G \implies M \#> \mathbf{1} = M$

$\langle proof \rangle$

lemma (**in group**) *coset-mult-inv1*:

$\llbracket M \#> (x \otimes (inv\ y)) = M; x \in carrier\ G; y \in carrier\ G; M \subseteq carrier\ G \rrbracket \implies M \#> x = M \#> y$

$\langle proof \rangle$

lemma (**in group**) *coset-mult-inv2*:

$\llbracket M \#> x = M \#> y; x \in carrier\ G; y \in carrier\ G; M \subseteq carrier\ G \rrbracket \implies M \#> (x \otimes (inv\ y)) = M$

$\langle proof \rangle$

lemma (**in group**) *coset-join1*:

$\llbracket H \#> x = H; x \in carrier\ G; subgroup\ H\ G \rrbracket \implies x \in H$

$\langle proof \rangle$

lemma (**in group**) *solve-equation*:

$\llbracket subgroup\ H\ G; x \in H; y \in H \rrbracket \implies \exists h \in H. y = h \otimes x$

$\langle proof \rangle$

lemma (**in group**) *repr-independence*:

$\llbracket y \in H \#> x; x \in carrier\ G; subgroup\ H\ G \rrbracket \implies H \#> x = H \#> y$

$\langle proof \rangle$

lemma (**in group**) *coset-join2*:

$\llbracket x \in carrier\ G; subgroup\ H\ G; x \in H \rrbracket \implies H \#> x = H$

— Alternative proof is to put $x = \mathbf{1}$ in *repr-independence*.

$\langle proof \rangle$

lemma (**in monoid**) *r-coset-subset-G*:

$\llbracket H \subseteq carrier\ G; x \in carrier\ G \rrbracket \implies H \#> x \subseteq carrier\ G$

$\langle proof \rangle$

lemma (**in group**) *rcosI*:

$\llbracket h \in H; H \subseteq carrier\ G; x \in carrier\ G \rrbracket \implies h \otimes x \in H \#> x$

$\langle proof \rangle$

lemma (in group) *rcosetsI*:

$\llbracket H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies H \#> x \in \text{rcosets } H$
 <proof>

Really needed?

lemma (in group) *transpose-inv*:

$\llbracket x \otimes y = z; x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket$
 $\implies (\text{inv } x) \otimes z = y$
 <proof>

lemma (in group) *rcos-self*: $\llbracket x \in \text{carrier } G; \text{subgroup } H \ G \rrbracket \implies x \in H \#> x$

<proof>

Opposite of *repr-independence*

lemma (in group) *repr-independenceD*:

includes *subgroup* $H \ G$
assumes *ycarr*: $y \in \text{carrier } G$
and *repr*: $H \#> x = H \#> y$
shows $y \in H \#> x$
 <proof>

Elements of a right coset are in the carrier

lemma (in subgroup) *elemrcos-carrier*:

includes *group*
assumes *acarr*: $a \in \text{carrier } G$
and *a'*: $a' \in H \#> a$
shows $a' \in \text{carrier } G$
 <proof>

lemma (in subgroup) *rcos-const*:

includes *group*
assumes *hH*: $h \in H$
shows $H \#> h = H$
 <proof>

Step one for lemma *rcos-module*

lemma (in subgroup) *rcos-module-imp*:

includes *group*
assumes *xcarr*: $x \in \text{carrier } G$
and *x'cos*: $x' \in H \#> x$
shows $(x' \otimes \text{inv } x) \in H$
 <proof>

Step two for lemma *rcos-module*

lemma (in subgroup) *rcos-module-rev*:

includes *group*
assumes *carr*: $x \in \text{carrier } G \ x' \in \text{carrier } G$
and *xiH*: $(x' \otimes \text{inv } x) \in H$

shows $x' \in H \#> x$
 ⟨proof⟩

Module property of right cosets

lemma (in *subgroup*) *rcos-module*:
includes *group*
assumes *carr*: $x \in \text{carrier } G$ $x' \in \text{carrier } G$
shows $(x' \in H \#> x) = (x' \otimes \text{inv } x \in H)$
 ⟨proof⟩

Right cosets are subsets of the carrier.

lemma (in *subgroup*) *rcosets-carrier*:
includes *group*
assumes *XH*: $X \in \text{rcosets } H$
shows $X \subseteq \text{carrier } G$
 ⟨proof⟩

Multiplication of general subsets

lemma (in *monoid*) *set-mult-closed*:
assumes *Acarr*: $A \subseteq \text{carrier } G$
and *Bcarr*: $B \subseteq \text{carrier } G$
shows $A <\#\> B \subseteq \text{carrier } G$
 ⟨proof⟩

lemma (in *comm-group*) *mult-subgroups*:
assumes *subH*: *subgroup* H G
and *subK*: *subgroup* K G
shows *subgroup* $(H <\#\> K)$ G
 ⟨proof⟩

lemma (in *subgroup*) *lcos-module-rev*:
includes *group*
assumes *carr*: $x \in \text{carrier } G$ $x' \in \text{carrier } G$
and *xixH*: $(\text{inv } x \otimes x') \in H$
shows $x' \in x <\#\> H$
 ⟨proof⟩

5.2 Normal subgroups

lemma *normal-imp-subgroup*: $H \triangleleft G \implies \text{subgroup } H$ G
 ⟨proof⟩

lemma (in *group*) *normalI*:
 $\text{subgroup } H$ $G \implies (\forall x \in \text{carrier } G. H \#> x = x <\#\> H) \implies H \triangleleft G$
 ⟨proof⟩

lemma (in *normal*) *inv-op-closed1*:
 $\llbracket x \in \text{carrier } G; h \in H \rrbracket \implies (\text{inv } x) \otimes h \otimes x \in H$
 ⟨proof⟩

lemma (in normal) *inv-op-closed2*:

$\llbracket x \in \text{carrier } G; h \in H \rrbracket \implies x \otimes h \otimes (\text{inv } x) \in H$
 <proof>

Alternative characterization of normal subgroups

lemma (in group) *normal-inv-iff*:

$(N \triangleleft G) =$
 (subgroup $N \ G$ & $(\forall x \in \text{carrier } G. \forall h \in N. x \otimes h \otimes (\text{inv } x) \in N)$)
 (is - = ?rhs)
 <proof>

5.3 More Properties of Cosets

lemma (in group) *lcos-m-assoc*:

$\llbracket M \subseteq \text{carrier } G; g \in \text{carrier } G; h \in \text{carrier } G \rrbracket$
 $\implies g <\# (h <\# M) = (g \otimes h) <\# M$
 <proof>

lemma (in group) *lcos-mult-one*: $M \subseteq \text{carrier } G \implies \mathbf{1} <\# M = M$

<proof>

lemma (in group) *l-coset-subset-G*:

$\llbracket H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies x <\# H \subseteq \text{carrier } G$
 <proof>

lemma (in group) *l-coset-swap*:

$\llbracket y \in x <\# H; x \in \text{carrier } G; \text{subgroup } H \ G \rrbracket \implies x \in y <\# H$
 <proof>

lemma (in group) *l-coset-carrier*:

$\llbracket y \in x <\# H; x \in \text{carrier } G; \text{subgroup } H \ G \rrbracket \implies y \in \text{carrier } G$
 <proof>

lemma (in group) *l-repr-imp-subset*:

assumes $y: y \in x <\# H$ and $x: x \in \text{carrier } G$ and $sb: \text{subgroup } H \ G$
 shows $y <\# H \subseteq x <\# H$
 <proof>

lemma (in group) *l-repr-independence*:

assumes $y: y \in x <\# H$ and $x: x \in \text{carrier } G$ and $sb: \text{subgroup } H \ G$
 shows $x <\# H = y <\# H$
 <proof>

lemma (in group) *setmult-subset-G*:

$\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G \rrbracket \implies H <\#> K \subseteq \text{carrier } G$
 <proof>

lemma (in group) *subgroup-mult-id*: $\text{subgroup } H \ G \implies H <\#> H = H$

<proof>

5.3.1 Set of Inverses of an r -coset.

lemma (in *normal*) *rcos-inv*:

assumes $x: x \in \text{carrier } G$

shows $\text{set-inv } (H \#> x) = H \#> (\text{inv } x)$

<proof>

5.3.2 Theorems for $<\#\>$ with $\#>$ or $<\#$.

lemma (in *group*) *setmult-rcos-assoc*:

$\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket$

$\implies H <\#\> (K \#> x) = (H <\#\> K) \#> x$

<proof>

lemma (in *group*) *rcos-assoc-lcos*:

$\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket$

$\implies (H \#> x) <\#\> K = H <\#\> (x <\# K)$

<proof>

lemma (in *normal*) *rcos-mult-step1*:

$\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket$

$\implies (H \#> x) <\#\> (H \#> y) = (H <\#\> (x <\# H)) \#> y$

<proof>

lemma (in *normal*) *rcos-mult-step2*:

$\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket$

$\implies (H <\#\> (x <\# H)) \#> y = (H <\#\> (H \#> x)) \#> y$

<proof>

lemma (in *normal*) *rcos-mult-step3*:

$\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket$

$\implies (H <\#\> (H \#> x)) \#> y = H \#> (x \otimes y)$

<proof>

lemma (in *normal*) *rcos-sum*:

$\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket$

$\implies (H \#> x) <\#\> (H \#> y) = H \#> (x \otimes y)$

<proof>

lemma (in *normal*) *rcosets-mult-eq*: $M \in \text{rcosets } H \implies H <\#\> M = M$

— generalizes *subgroup-mult-id*

<proof>

5.3.3 An Equivalence Relation

constdefs (structure G)

$r\text{-congruent} :: [(\text{'a}, \text{'b}) \text{monoid-scheme}, \text{'a set}] \Rightarrow (\text{'a} * \text{'a}) \text{set}$
 $(\text{rcong1 } -)$

$rcong\ H \equiv \{(x,y). x \in carrier\ G \ \& \ y \in carrier\ G \ \& \ inv\ x \otimes y \in H\}$

lemma (in *subgroup*) *equiv-rcong*:
includes *group* G
shows *equiv* (*carrier* G) (*rcong* H)
 ⟨*proof*⟩

Equivalence classes of *rcong* correspond to left cosets. Was there a mistake in the definitions? I'd have expected them to correspond to right cosets.

lemma (in *subgroup*) *l-coset-eq-rcong*:
includes *group* G
assumes $a: a \in carrier\ G$
shows $a <\# H = rcong\ H\ \{\{a\}\}$
 ⟨*proof*⟩

5.3.4 Two Distinct Right Cosets are Disjoint

lemma (in *group*) *rcos-equation*:
includes *subgroup* $H\ G$
shows
 $\llbracket ha \otimes a = h \otimes b; a \in carrier\ G; b \in carrier\ G;$
 $h \in H; ha \in H; hb \in H \rrbracket$
 $\implies hb \otimes a \in (\bigcup h \in H. \{h \otimes b\})$
 ⟨*proof*⟩

lemma (in *group*) *rcos-disjoint*:
includes *subgroup* $H\ G$
shows $\llbracket a \in rcosets\ H; b \in rcosets\ H; a \neq b \rrbracket \implies a \cap b = \{\}$
 ⟨*proof*⟩

5.4 Further lemmas for *r-congruent*

The relation is a congruence

lemma (in *normal*) *congruent-rcong*:
shows *congruent2* (*rcong* H) (*rcong* H) ($\lambda a\ b. a \otimes b <\# H$)
 ⟨*proof*⟩

5.5 Order of a Group and Lagrange's Theorem

constdefs
 $order :: ('a, 'b)\ monoid\ scheme \Rightarrow nat$
 $order\ S \equiv card\ (carrier\ S)$

lemma (in *group*) *rcos-self*:
includes *subgroup*
shows $x \in carrier\ G \implies x \in H \ \#> x$
 ⟨*proof*⟩

lemma (in group) *rcosets-part-G*:
includes *subgroup*
shows $\bigcup (\text{rcosets } H) = \text{carrier } G$
 ⟨proof⟩

lemma (in group) *cosets-finite*:
 $\llbracket c \in \text{rcosets } H; H \subseteq \text{carrier } G; \text{finite } (\text{carrier } G) \rrbracket \implies \text{finite } c$
 ⟨proof⟩

The next two lemmas support the proof of *card-cosets-equal*.

lemma (in group) *inj-on-f*:
 $\llbracket H \subseteq \text{carrier } G; a \in \text{carrier } G \rrbracket \implies \text{inj-on } (\lambda y. y \otimes \text{inv } a) (H \#> a)$
 ⟨proof⟩

lemma (in group) *inj-on-g*:
 $\llbracket H \subseteq \text{carrier } G; a \in \text{carrier } G \rrbracket \implies \text{inj-on } (\lambda y. y \otimes a) H$
 ⟨proof⟩

lemma (in group) *card-cosets-equal*:
 $\llbracket c \in \text{rcosets } H; H \subseteq \text{carrier } G; \text{finite}(\text{carrier } G) \rrbracket$
 $\implies \text{card } c = \text{card } H$
 ⟨proof⟩

lemma (in group) *rcosets-subset-PowG*:
 $\text{subgroup } H \ G \implies \text{rcosets } H \subseteq \text{Pow}(\text{carrier } G)$
 ⟨proof⟩

theorem (in group) *lagrange*:
 $\llbracket \text{finite}(\text{carrier } G); \text{subgroup } H \ G \rrbracket$
 $\implies \text{card}(\text{rcosets } H) * \text{card}(H) = \text{order}(G)$
 ⟨proof⟩

5.6 Quotient Groups: Factorization of a Group

constdefs

FactGroup :: $[('a, 'b) \text{ monoid-scheme}, 'a \text{ set}] \Rightarrow ('a \text{ set}) \text{ monoid}$
 (infixl Mod 65)
 — Actually defined for groups rather than monoids
 $\text{FactGroup } G \ H \equiv$
 $(\text{carrier} = \text{rcosets } G \ H, \text{mult} = \text{set-mult } G, \text{one} = H)$

lemma (in normal) *setmult-closed*:
 $\llbracket K1 \in \text{rcosets } H; K2 \in \text{rcosets } H \rrbracket \implies K1 \langle \# \rangle K2 \in \text{rcosets } H$
 ⟨proof⟩

lemma (in normal) *setinv-closed*:
 $K \in \text{rcosets } H \implies \text{set-inv } K \in \text{rcosets } H$
 ⟨proof⟩

lemma (in normal) *rcosets-assoc*:

$\llbracket M1 \in \text{rcosets } H; M2 \in \text{rcosets } H; M3 \in \text{rcosets } H \rrbracket$
 $\implies M1 <\#\rangle M2 <\#\rangle M3 = M1 <\#\rangle (M2 <\#\rangle M3)$
 ⟨proof⟩

lemma (in subgroup) *subgroup-in-rcosets*:

includes group G
shows $H \in \text{rcosets } H$
 ⟨proof⟩

lemma (in normal) *rcosets-inv-mult-group-eq*:

$M \in \text{rcosets } H \implies \text{set-inv } M <\#\rangle M = H$
 ⟨proof⟩

theorem (in normal) *factorgroup-is-group*:

group $(G \text{ Mod } H)$
 ⟨proof⟩

lemma *mult-FactGroup [simp]*: $X \otimes_{(G \text{ Mod } H)} X' = X <\#\rangle_G X'$

⟨proof⟩

lemma (in normal) *inv-FactGroup*:

$X \in \text{carrier } (G \text{ Mod } H) \implies \text{inv}_{G \text{ Mod } H} X = \text{set-inv } X$
 ⟨proof⟩

The coset map is a homomorphism from G to the quotient group $G \text{ Mod } H$

lemma (in normal) *r-coset-hom-Mod*:

$(\lambda a. H \#\rangle a) \in \text{hom } G (G \text{ Mod } H)$
 ⟨proof⟩

5.7 The First Isomorphism Theorem

The quotient by the kernel of a homomorphism is isomorphic to the range of that homomorphism.

constdefs

kernel :: ('a, 'm) monoid-scheme \Rightarrow ('b, 'n) monoid-scheme \Rightarrow
 ('a \Rightarrow 'b) \Rightarrow 'a set
 — the kernel of a homomorphism
 $\text{kernel } G H h \equiv \{x. x \in \text{carrier } G \ \& \ h \ x = \mathbf{1}_H\}$

lemma (in group-hom) *subgroup-kernel*: *subgroup* (kernel $G H h$) G

⟨proof⟩

The kernel of a homomorphism is a normal subgroup

lemma (in group-hom) *normal-kernel*: (kernel $G H h$) $\triangleleft G$

⟨proof⟩

lemma (in *group-hom*) *FactGroup-nonempty*:
 assumes $X: X \in \text{carrier } (G \text{ Mod } \text{kernel } G \ H \ h)$
 shows $X \neq \{\}$
 <proof>

lemma (in *group-hom*) *FactGroup-contents-mem*:
 assumes $X: X \in \text{carrier } (G \text{ Mod } (\text{kernel } G \ H \ h))$
 shows $\text{contents } (h'X) \in \text{carrier } H$
 <proof>

lemma (in *group-hom*) *FactGroup-hom*:
 $(\lambda X. \text{contents } (h'X)) \in \text{hom } (G \text{ Mod } (\text{kernel } G \ H \ h)) \ H$
 <proof>

Lemma for the following injectivity result

lemma (in *group-hom*) *FactGroup-subset*:
 $\llbracket g \in \text{carrier } G; g' \in \text{carrier } G; h \ g = h \ g' \rrbracket$
 $\implies \text{kernel } G \ H \ h \ \#> \ g \subseteq \text{kernel } G \ H \ h \ \#> \ g'$
 <proof>

lemma (in *group-hom*) *FactGroup-inj-on*:
 $\text{inj-on } (\lambda X. \text{contents } (h'X)) \ (\text{carrier } (G \text{ Mod } \text{kernel } G \ H \ h))$
 <proof>

If the homomorphism h is onto H , then so is the homomorphism from the quotient group

lemma (in *group-hom*) *FactGroup-onto*:
 assumes $h: h' \text{carrier } G = \text{carrier } H$
 shows $(\lambda X. \text{contents } (h'X))' \text{carrier } (G \text{ Mod } \text{kernel } G \ H \ h) = \text{carrier } H$
 <proof>

If h is a homomorphism from G onto H , then the quotient group $G \text{ Mod } \text{kernel } G \ H \ h$ is isomorphic to H .

theorem (in *group-hom*) *FactGroup-iso*:
 $h' \text{carrier } G = \text{carrier } H$
 $\implies (\lambda X. \text{contents } (h'X)) \in (G \text{ Mod } (\text{kernel } G \ H \ h)) \cong H$
 <proof>

end

theory *Sylow* imports *Coset* begin

6 Sylow's Theorem

See also [3].

The combinatorial argument is in theory Exponent

locale *sylow* = *group* +
fixes *p* **and** *a* **and** *m* **and** *calM* **and** *RelM*
assumes *prime-p*: *prime p*
and *order-G*: $order(G) = (p \wedge a) * m$
and *finite-G* [*iff*]: *finite (carrier G)*
defines *calM* == $\{s. s \subseteq carrier(G) \ \& \ card(s) = p \wedge a\}$
and *RelM* == $\{(N1, N2). N1 \in calM \ \& \ N2 \in calM \ \& \ (\exists g \in carrier(G). N1 = (N2 \#> g))\}$

lemma (**in** *sylow*) *RelM-refl*: *refl calM RelM*
<proof>

lemma (**in** *sylow*) *RelM-sym*: *sym RelM*
<proof>

lemma (**in** *sylow*) *RelM-trans*: *trans RelM*
<proof>

lemma (**in** *sylow*) *RelM-equiv*: *equiv calM RelM*
<proof>

lemma (**in** *sylow*) *M-subset-calM-prep*: $M' \in calM // RelM ==> M' \subseteq calM$
<proof>

6.1 Main Part of the Proof

locale *sylow-central* = *sylow* +
fixes *H* **and** *M1* **and** *M*
assumes *M-in-quot*: $M \in calM // RelM$
and *not-dvd-M*: $\sim(p \wedge Suc(exponent\ p\ m) \ dvd\ card(M))$
and *M1-in-M*: $M1 \in M$
defines *H* == $\{g. g \in carrier\ G \ \& \ M1 \ \#> \ g = M1\}$

lemma (**in** *sylow-central*) *M-subset-calM*: $M \subseteq calM$
<proof>

lemma (**in** *sylow-central*) *card-M1*: $card(M1) = p \wedge a$
<proof>

lemma *card-nonempty*: $0 < card(S) ==> S \neq \{\}$
<proof>

lemma (**in** *sylow-central*) *exists-x-in-M1*: $\exists x. x \in M1$
<proof>

lemma (in *syLOW-central*) *M1-subset-G* [*simp*]: $M1 \subseteq \text{carrier } G$
 ⟨*proof*⟩

lemma (in *syLOW-central*) *M1-inj-H*: $\exists f \in H \rightarrow M1. \text{inj-on } f \ H$
 ⟨*proof*⟩

6.2 Discharging the Assumptions of *syLOW-central*

lemma (in *syLOW*) *EmptyNotInEquivSet*: $\{\} \notin \text{calM} // \text{RelM}$
 ⟨*proof*⟩

lemma (in *syLOW*) *existsM1inM*: $M \in \text{calM} // \text{RelM} \implies \exists M1. M1 \in M$
 ⟨*proof*⟩

lemma (in *syLOW*) *zero-less-o-G*: $0 < \text{order}(G)$
 ⟨*proof*⟩

lemma (in *syLOW*) *zero-less-m*: $m > 0$
 ⟨*proof*⟩

lemma (in *syLOW*) *card-calM*: $\text{card}(\text{calM}) = (p^a) * m$ choose p^a
 ⟨*proof*⟩

lemma (in *syLOW*) *zero-less-card-calM*: $\text{card } \text{calM} > 0$
 ⟨*proof*⟩

lemma (in *syLOW*) *max-p-div-calM*:
 $\sim (p \wedge \text{Suc}(\text{exponent } p \ m) \ \text{dvd } \text{card}(\text{calM}))$
 ⟨*proof*⟩

lemma (in *syLOW*) *finite-calM*: *finite calM*
 ⟨*proof*⟩

lemma (in *syLOW*) *lemma-A1*:
 $\exists M \in \text{calM} // \text{RelM}. \sim (p \wedge \text{Suc}(\text{exponent } p \ m) \ \text{dvd } \text{card}(M))$
 ⟨*proof*⟩

6.2.1 Introduction and Destruct Rules for *H*

lemma (in *syLOW-central*) *H-I*: $[[g \in \text{carrier } G; M1 \#> g = M1]] \implies g \in H$
 ⟨*proof*⟩

lemma (in *syLOW-central*) *H-into-carrier-G*: $x \in H \implies x \in \text{carrier } G$
 ⟨*proof*⟩

lemma (in *syLOW-central*) *in-H-imp-eq*: $g : H \implies M1 \#> g = M1$
 ⟨*proof*⟩

lemma (in *syLOW-central*) *H-m-closed*: $[[x \in H; y \in H]] \implies x \otimes y \in H$

<proof>

lemma (in *syLOW-central*) *H-not-empty*: $H \neq \{\}$
<proof>

lemma (in *syLOW-central*) *H-is-subgroup*: subgroup $H \ G$
<proof>

lemma (in *syLOW-central*) *rcosetGM1g-subset-G*:
 $[\![\ g \in \text{carrier } G; x \in M1 \ \#> \ g \]\!] \implies x \in \text{carrier } G$
<proof>

lemma (in *syLOW-central*) *finite-M1*: finite $M1$
<proof>

lemma (in *syLOW-central*) *finite-rcosetGM1g*: $g \in \text{carrier } G \implies \text{finite } (M1 \ \#> \ g)$
<proof>

lemma (in *syLOW-central*) *M1-card-eq-rcosetGM1g*:
 $g \in \text{carrier } G \implies \text{card}(M1 \ \#> \ g) = \text{card}(M1)$
<proof>

lemma (in *syLOW-central*) *M1-RelM-rcosetGM1g*:
 $g \in \text{carrier } G \implies (M1, M1 \ \#> \ g) \in \text{RelM}$
<proof>

6.3 Equal Cardinalities of M and the Set of Cosets

Injections between M and $\text{rcosets}_G H$ show that their cardinalities are equal.

lemma *ElemClassEquiv*:
 $[\![\ \text{equiv } A \ r; C \in A \ // \ r \]\!] \implies \forall x \in C. \forall y \in C. (x,y) \in r$
<proof>

lemma (in *syLOW-central*) *M-elem-map*:
 $M2 \in M \implies \exists g. g \in \text{carrier } G \ \& \ M1 \ \#> \ g = M2$
<proof>

lemmas (in *syLOW-central*) *M-elem-map-carrier =*
 $M\text{-elem-map } [\text{THEN someI-ex}, \text{THEN conjunct1}]$

lemmas (in *syLOW-central*) *M-elem-map-eq =*
 $M\text{-elem-map } [\text{THEN someI-ex}, \text{THEN conjunct2}]$

lemma (in *syLOW-central*) *M-funcset-rcosets-H*:
 $(\%x:M. H \ \#> \ (\text{SOME } g. g \in \text{carrier } G \ \& \ M1 \ \#> \ g = x)) \in M \rightarrow \text{rcosets } H$
<proof>

lemma (in *syLOW-central*) *inj-M-GmodH*: $\exists f \in M \rightarrow \text{rcosets } H. \text{inj-on } f \ M$

<proof>

6.3.1 The Opposite Injection

lemma (in *syLOW-central*) *H-elem-map*:

$$H1 \in \text{rcosets } H \implies \exists g. g \in \text{carrier } G \ \& \ H \ \#> \ g = H1$$

<proof>

lemmas (in *syLOW-central*) *H-elem-map-carrier =*

$$H\text{-elem-map [THEN someI-ex, THEN conjunct1]}$$

lemmas (in *syLOW-central*) *H-elem-map-eq =*

$$H\text{-elem-map [THEN someI-ex, THEN conjunct2]}$$

lemma *EquivElemClass*:

$$[[\text{equiv } A \ r; \ M \in A//r; \ M1 \in M; \ (M1, M2) \in r \] \implies M2 \in M$$

<proof>

lemma (in *syLOW-central*) *rcosets-H-funcset-M*:

$$(\lambda C \in \text{rcosets } H. M1 \ \#> \ (@g. g \in \text{carrier } G \ \wedge \ H \ \#> \ g = C)) \in \text{rcosets } H \ \rightarrow M$$

<proof>

close to a duplicate of *inj-M-GmodH*

lemma (in *syLOW-central*) *inj-GmodH-M*:

$$\exists g \in \text{rcosets } H \rightarrow M. \text{inj-on } g \ (\text{rcosets } H)$$

<proof>

lemma (in *syLOW-central*) *calM-subset-PowG*: $\text{cal}M \subseteq \text{Pow}(\text{carrier } G)$

<proof>

lemma (in *syLOW-central*) *finite-M*: *finite* M

<proof>

lemma (in *syLOW-central*) *cardMeqIndexH*: $\text{card}(M) = \text{card}(\text{rcosets } H)$

<proof>

lemma (in *syLOW-central*) *index-lem*: $\text{card}(M) * \text{card}(H) = \text{order}(G)$

<proof>

lemma (in *syLOW-central*) *lemma-leq1*: $p^a \leq \text{card}(H)$

<proof>

lemma (in *syLOW-central*) *lemma-leq2*: $\text{card}(H) \leq p^a$

<proof>

lemma (in *syLOW-central*) *card-H-eq*: $\text{card}(H) = p^a$
 ⟨*proof*⟩

lemma (in *syLOW*) *syLOW-thm*: $\exists H. \text{subgroup } H \ G \ \& \ \text{card}(H) = p^a$
 ⟨*proof*⟩

Needed because the locale's automatic definition refers to *semigroup* G and *group-axioms* G rather than simply to *group* G .

lemma *syLOW-eq*: $\text{syLOW } G \ p \ a \ m = (\text{group } G \ \& \ \text{syLOW-axioms } G \ p \ a \ m)$
 ⟨*proof*⟩

6.4 Sylow's Theorem

theorem *syLOW-thm*:

[[*prime* p ; *group*(G); $\text{order}(G) = (p^a) * m$; *finite* (*carrier* G)]
 $\implies \exists H. \text{subgroup } H \ G \ \& \ \text{card}(H) = p^a$

⟨*proof*⟩

end

theory *Bij* imports *Group* begin

7 Bijections of a Set, Permutation Groups and Automorphism Groups

constdefs

Bij :: $'a \text{ set} \Rightarrow ('a \Rightarrow 'a) \text{ set}$

— Only extensional functions, since otherwise we get too many.

$\text{Bij } S \equiv \text{extensional } S \cap \{f. \text{bij-betw } f \ S \ S\}$

BijGroup :: $'a \text{ set} \Rightarrow ('a \Rightarrow 'a) \text{ monoid}$

$\text{BijGroup } S \equiv$

($\text{carrier} = \text{Bij } S$,

$\text{mult} = \lambda g \in \text{Bij } S. \lambda f \in \text{Bij } S. \text{compose } S \ g \ f$,

$\text{one} = \lambda x \in S. x$)

declare *Id-compose* [*simp*] *compose-Id* [*simp*]

lemma *Bij-imp-extensional*: $f \in \text{Bij } S \implies f \in \text{extensional } S$
 ⟨*proof*⟩

lemma *Bij-imp-funcset*: $f \in \text{Bij } S \implies f \in S \rightarrow S$
 ⟨*proof*⟩

7.1 Bijections Form a Group

lemma *restrict-Inv-Bij*: $f \in \text{Bij } S \implies (\lambda x \in S. (\text{Inv } S f) x) \in \text{Bij } S$
 ⟨proof⟩

lemma *id-Bij*: $(\lambda x \in S. x) \in \text{Bij } S$
 ⟨proof⟩

lemma *compose-Bij*: $\llbracket x \in \text{Bij } S; y \in \text{Bij } S \rrbracket \implies \text{compose } S x y \in \text{Bij } S$
 ⟨proof⟩

lemma *Bij-compose-restrict-eq*:
 $f \in \text{Bij } S \implies \text{compose } S (\text{restrict } (\text{Inv } S f) S) f = (\lambda x \in S. x)$
 ⟨proof⟩

theorem *group-BijGroup*: $\text{group } (\text{BijGroup } S)$
 ⟨proof⟩

7.2 Automorphisms Form a Group

lemma *Bij-Inv-mem*: $\llbracket f \in \text{Bij } S; x \in S \rrbracket \implies \text{Inv } S f x \in S$
 ⟨proof⟩

lemma *Bij-Inv-lemma*:
assumes *eq*: $\bigwedge x y. \llbracket x \in S; y \in S \rrbracket \implies h(g x y) = g (h x) (h y)$
shows $\llbracket h \in \text{Bij } S; g \in S \rightarrow S \rightarrow S; x \in S; y \in S \rrbracket$
 $\implies \text{Inv } S h (g x y) = g (\text{Inv } S h x) (\text{Inv } S h y)$
 ⟨proof⟩

constdefs

auto :: $('a, 'b) \text{ monoid-scheme} \Rightarrow ('a \Rightarrow 'a) \text{ set}$
auto $G \equiv \text{hom } G G \cap \text{Bij } (\text{carrier } G)$

AutoGroup :: $('a, 'c) \text{ monoid-scheme} \Rightarrow ('a \Rightarrow 'a) \text{ monoid}$
AutoGroup $G \equiv \text{BijGroup } (\text{carrier } G) (\downarrow \text{carrier} := \text{auto } G)$

lemma (in *group*) *id-in-auto*: $(\lambda x \in \text{carrier } G. x) \in \text{auto } G$
 ⟨proof⟩

lemma (in *group*) *mult-funcset*: $\text{mult } G \in \text{carrier } G \rightarrow \text{carrier } G \rightarrow \text{carrier } G$
 ⟨proof⟩

lemma (in *group*) *restrict-Inv-hom*:
 $\llbracket h \in \text{hom } G G; h \in \text{Bij } (\text{carrier } G) \rrbracket$
 $\implies \text{restrict } (\text{Inv } (\text{carrier } G) h) (\text{carrier } G) \in \text{hom } G G$
 ⟨proof⟩

lemma *inv-BijGroup*:
 $f \in \text{Bij } S \implies \text{m-inv } (\text{BijGroup } S) f = (\lambda x \in S. (\text{Inv } S f) x)$

<proof>

lemma (*in group*) *subgroup-auto*:
subgroup (auto G) (BijGroup (carrier G))
<proof>

theorem (*in group*) *AutoGroup: group (AutoGroup G)*
<proof>

end

theory *Ring imports FiniteProduct*
uses (*ringsimp.ML*) **begin**

8 Abelian Groups

record *'a ring = 'a monoid +*
zero :: 'a (01)
add :: ['a, 'a] => 'a (infixl \oplus 65)

Derived operations.

constdefs (**structure** *R*)
a-inv :: [('a, 'm) ring-scheme, 'a] => 'a (\ominus 1 - [81] 80)
a-inv R == m-inv (| carrier = carrier R, mult = add R, one = zero R |)

a-minus :: [('a, 'm) ring-scheme, 'a, 'a] => 'a (infixl \ominus 1 65)
[| x \in carrier R; y \in carrier R |] ==> x \ominus y == x \oplus (\ominus y)

locale *abelian-monoid =*
fixes *G (structure)*
assumes *a-comm-monoid*:
comm-monoid (| carrier = carrier G, mult = add G, one = zero G |)

The following definition is redundant but simple to use.

locale *abelian-group = abelian-monoid +*
assumes *a-comm-group*:
comm-group (| carrier = carrier G, mult = add G, one = zero G |)

8.1 Basic Properties

lemma *abelian-monoidI*:
fixes *R (structure)*
assumes *a-closed*:
!!x y. [| x \in carrier R; y \in carrier R |] ==> x \oplus y \in carrier R
and *zero-closed*: *0 \in carrier R*
and *a-assoc*:

$!!x\ y\ z. [\![\ x \in \text{carrier } R; y \in \text{carrier } R; z \in \text{carrier } R \]\!] ==>$
 $(x \oplus y) \oplus z = x \oplus (y \oplus z)$
and *l-zero*: $!!x. x \in \text{carrier } R ==> \mathbf{0} \oplus x = x$
and *a-comm*:
 $!!x\ y. [\![\ x \in \text{carrier } R; y \in \text{carrier } R \]\!] ==> x \oplus y = y \oplus x$
shows *abelian-monoid* R
 $\langle \text{proof} \rangle$

lemma *abelian-groupI*:
fixes R (**structure**)
assumes *a-closed*:
 $!!x\ y. [\![\ x \in \text{carrier } R; y \in \text{carrier } R \]\!] ==> x \oplus y \in \text{carrier } R$
and *zero-closed*: $\text{zero } R \in \text{carrier } R$
and *a-assoc*:
 $!!x\ y\ z. [\![\ x \in \text{carrier } R; y \in \text{carrier } R; z \in \text{carrier } R \]\!] ==>$
 $(x \oplus y) \oplus z = x \oplus (y \oplus z)$
and *a-comm*:
 $!!x\ y. [\![\ x \in \text{carrier } R; y \in \text{carrier } R \]\!] ==> x \oplus y = y \oplus x$
and *l-zero*: $!!x. x \in \text{carrier } R ==> \mathbf{0} \oplus x = x$
and *l-inv-ex*: $!!x. x \in \text{carrier } R ==> \exists y : \text{carrier } R. y \oplus x = \mathbf{0}$
shows *abelian-group* R
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-monoid*) *a-monoid*:
 $\text{monoid } (\mid \text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G \mid)$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-group*) *a-group*:
 $\text{group } (\mid \text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G \mid)$
 $\langle \text{proof} \rangle$

lemmas *monoid-record-simps* = *partial-object.simps monoid.simps*

lemma (**in** *abelian-monoid*) *a-closed* [*intro*, *simp*]:
 $[\![\ x \in \text{carrier } G; y \in \text{carrier } G \]\!] ==> x \oplus y \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-monoid*) *zero-closed* [*intro*, *simp*]:
 $\mathbf{0} \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-group*) *a-inv-closed* [*intro*, *simp*]:
 $x \in \text{carrier } G ==> \ominus x \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-group*) *minus-closed* [*intro*, *simp*]:
 $[\![\ x \in \text{carrier } G; y \in \text{carrier } G \]\!] ==> x \ominus y \in \text{carrier } G$
 $\langle \text{proof} \rangle$

lemma (*in abelian-group*) *a-l-cancel* [*simp*]:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$
 $(x \oplus y = x \oplus z) = (y = z)$
<proof>

lemma (*in abelian-group*) *a-r-cancel* [*simp*]:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$
 $(y \oplus x = z \oplus x) = (y = z)$
<proof>

lemma (*in abelian-monoid*) *a-assoc*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$
 $(x \oplus y) \oplus z = x \oplus (y \oplus z)$
<proof>

lemma (*in abelian-monoid*) *l-zero* [*simp*]:
 $x \in \text{carrier } G \implies \mathbf{0} \oplus x = x$
<proof>

lemma (*in abelian-group*) *l-neg*:
 $x \in \text{carrier } G \implies \ominus x \oplus x = \mathbf{0}$
<proof>

lemma (*in abelian-monoid*) *a-comm*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies x \oplus y = y \oplus x$
<proof>

lemma (*in abelian-monoid*) *a-lcomm*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket \implies$
 $x \oplus (y \oplus z) = y \oplus (x \oplus z)$
<proof>

lemma (*in abelian-monoid*) *r-zero* [*simp*]:
 $x \in \text{carrier } G \implies x \oplus \mathbf{0} = x$
<proof>

lemma (*in abelian-group*) *r-neg*:
 $x \in \text{carrier } G \implies x \oplus (\ominus x) = \mathbf{0}$
<proof>

lemma (*in abelian-group*) *minus-zero* [*simp*]:
 $\ominus \mathbf{0} = \mathbf{0}$
<proof>

lemma (*in abelian-group*) *minus-minus* [*simp*]:
 $x \in \text{carrier } G \implies \ominus (\ominus x) = x$
<proof>

lemma (*in abelian-group*) *a-inv-inj*:

inj-on (*a-inv* *G*) (*carrier* *G*)
 ⟨*proof*⟩

lemma (*in abelian-group*) *minus-add*:
 [| *x* ∈ *carrier* *G*; *y* ∈ *carrier* *G* |] ==> $\ominus (x \oplus y) = \ominus x \oplus \ominus y$
 ⟨*proof*⟩

lemma (*in abelian-group*) *minus-equality*:
 [| *x* ∈ *carrier* *G*; *y* ∈ *carrier* *G*; *y* ⊕ *x* = **0** |] ==> $\ominus x = y$
 ⟨*proof*⟩

lemma (*in abelian-monoid*) *minus-unique*:
 [| *x* ∈ *carrier* *G*; *y* ∈ *carrier* *G*; *y*' ∈ *carrier* *G*;
y ⊕ *x* = **0**; *x* ⊕ *y*' = **0** |] ==> *y* = *y*'
 ⟨*proof*⟩

lemmas (*in abelian-monoid*) *a-ac = a-assoc a-comm a-lcomm*

Derive an *abelian-group* from a *comm-group*

lemma *comm-group-abelian-groupI*:
fixes *G* (**structure**)
assumes *cg*: *comm-group* (| *carrier* = *carrier* *G*, *mult* = *add* *G*, *one* = *zero* *G*)
shows *abelian-group* *G*
 ⟨*proof*⟩

8.2 Sums over Finite Sets

This definition makes it easy to lift lemmas from *finprod*.

constdefs
finsum :: [(*'b*, *'m*) *ring-scheme*, *'a* => *'b*, *'a set*] => *'b*
finsum *G* *f* *A* == *finprod* (| *carrier* = *carrier* *G*,
mult = *add* *G*, *one* = *zero* *G* |) *f* *A*

syntax
 -*finsum* :: *index* => *idt* => *'a set* => *'b* => *'b*
 (($\exists \oplus$ --:.-) [1000, 0, 51, 10] 10)

syntax (*xsymbols*)
 -*finsum* :: *index* => *idt* => *'a set* => *'b* => *'b*
 (($\exists \oplus$ --∈.-) [1000, 0, 51, 10] 10)

syntax (*HTML output*)
 -*finsum* :: *index* => *idt* => *'a set* => *'b* => *'b*
 (($\exists \oplus$ --∈.-) [1000, 0, 51, 10] 10)

translations
 $\bigoplus_{i:A} b == finsum \circ_1 (\%i. b) A$
 — Beware of argument permutation!

lemma (*in abelian-monoid*) *finsum-empty* [*simp*]:

$\text{finsum } G f \{\} = \mathbf{0}$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-insert* [simp]:
 [| finite F; a ∉ F; f ∈ F -> carrier G; f a ∈ carrier G |]
 ==> $\text{finsum } G f (\text{insert } a F) = f a \oplus \text{finsum } G f F$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-zero* [simp]:
 finite A ==> $(\bigoplus_{i \in A} \mathbf{0}) = \mathbf{0}$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-closed* [simp]:
 fixes A
 assumes fin: finite A and f: f ∈ A -> carrier G
 shows $\text{finsum } G f A \in \text{carrier } G$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-Un-Int*:
 [| finite A; finite B; g ∈ A -> carrier G; g ∈ B -> carrier G |] ==>
 $\text{finsum } G g (A \text{ Un } B) \oplus \text{finsum } G g (A \text{ Int } B) =$
 $\text{finsum } G g A \oplus \text{finsum } G g B$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-Un-disjoint*:
 [| finite A; finite B; A Int B = {};
 g ∈ A -> carrier G; g ∈ B -> carrier G |]
 ==> $\text{finsum } G g (A \text{ Un } B) = \text{finsum } G g A \oplus \text{finsum } G g B$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-addr*:
 [| finite A; f ∈ A -> carrier G; g ∈ A -> carrier G |] ==>
 $\text{finsum } G (\%x. f x \oplus g x) A = (\text{finsum } G f A \oplus \text{finsum } G g A)$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-cong'*:
 [| A = B; g : B -> carrier G;
 !!i. i : B ==> f i = g i |] ==> $\text{finsum } G f A = \text{finsum } G g B$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-0* [simp]:
 f : {0::nat} -> carrier G ==> $\text{finsum } G f \{..0\} = f 0$
 ⟨proof⟩

lemma (in *abelian-monoid*) *finsum-Suc* [simp]:
 f : {..Suc n} -> carrier G ==>
 $\text{finsum } G f \{..Suc n\} = (f (\text{Suc } n) \oplus \text{finsum } G f \{..n\})$
 ⟨proof⟩

$==> z \otimes (x \oplus y) = z \otimes x \oplus z \otimes y$
shows *ring* R
 <proof>

lemma (**in** *ring*) *is-abelian-group*:
abelian-group R
 <proof>

lemma (**in** *ring*) *is-monoid*:
monoid R
 <proof>

lemma (**in** *ring*) *is-ring*:
ring R
 <proof>

lemmas *ring-record-simps* = *monoid-record-simps* *ring.simps*

lemma *cringI*:
fixes R (**structure**)
assumes *abelian-group*: *abelian-group* R
and *comm-monoid*: *comm-monoid* R
and *l-distr*: $!!x\ y\ z. [| x \in \text{carrier } R; y \in \text{carrier } R; z \in \text{carrier } R |]$
 $==> (x \oplus y) \otimes z = x \otimes z \oplus y \otimes z$
shows *cring* R
 <proof>

lemma (**in** *cring*) *is-comm-monoid*:
comm-monoid R
 <proof>

lemma (**in** *cring*) *is-cring*:
cring R <proof>

9.2.1 Normaliser for Rings

lemma (**in** *abelian-group*) *r-neg2*:
 $[| x \in \text{carrier } G; y \in \text{carrier } G |] ==> x \oplus (\ominus x \oplus y) = y$
 <proof>

lemma (**in** *abelian-group*) *r-neg1*:
 $[| x \in \text{carrier } G; y \in \text{carrier } G |] ==> \ominus x \oplus (x \oplus y) = y$
 <proof>

The following proofs are from Jacobson, Basic Algebra I, pp. 88–89

lemma (**in** *ring*) *l-null* [*simp*]:
 $x \in \text{carrier } R ==> \mathbf{0} \otimes x = \mathbf{0}$
 <proof>

lemma (in ring) *r-null [simp]*:
 $x \in \text{carrier } R \implies x \otimes \mathbf{0} = \mathbf{0}$
 <proof>

lemma (in ring) *l-minus*:
 $[[x \in \text{carrier } R; y \in \text{carrier } R]] \implies \ominus x \otimes y = \ominus (x \otimes y)$
 <proof>

lemma (in ring) *r-minus*:
 $[[x \in \text{carrier } R; y \in \text{carrier } R]] \implies x \otimes \ominus y = \ominus (x \otimes y)$
 <proof>

lemma (in abelian-group) *minus-eq*:
 $[[x \in \text{carrier } G; y \in \text{carrier } G]] \implies x \ominus y = x \oplus \ominus y$
 <proof>

Setup algebra method: compute distributive normal form in locale contexts

<ML>

lemmas (in ring) *ring-simprules*
 $[\text{algebra ring zero } R \text{ add } R \text{ a-inv } R \text{ a-minus } R \text{ one } R \text{ mult } R] =$
a-closed zero-closed a-inv-closed minus-closed m-closed one-closed
a-assoc l-zero l-neg a-comm m-assoc l-one l-distr minus-eq
r-zero r-neg r-neg2 r-neg1 minus-add minus-minus minus-zero
a-lcomm r-distr l-null r-null l-minus r-minus

lemmas (in cring)
 $[\text{algebra del: ring zero } R \text{ add } R \text{ a-inv } R \text{ a-minus } R \text{ one } R \text{ mult } R] =$
 -

lemmas (in cring) *cring-simprules*
 $[\text{algebra add: cring zero } R \text{ add } R \text{ a-inv } R \text{ a-minus } R \text{ one } R \text{ mult } R] =$
a-closed zero-closed a-inv-closed minus-closed m-closed one-closed
a-assoc l-zero l-neg a-comm m-assoc l-one l-distr m-comm minus-eq
r-zero r-neg r-neg2 r-neg1 minus-add minus-minus minus-zero
a-lcomm m-lcomm r-distr l-null r-null l-minus r-minus

lemma (in cring) *nat-pow-zero*:
 $(n::\text{nat}) \sim 0 \implies \mathbf{0} (\wedge) n = \mathbf{0}$
 <proof>

lemma (in ring) *one-zeroD*:
assumes *onezero*: $\mathbf{1} = \mathbf{0}$
shows *carrier* $R = \{\mathbf{0}\}$
 <proof>

lemma (in ring) *one-zeroI*:
assumes *carrzero*: *carrier* $R = \{\mathbf{0}\}$

shows $\mathbf{1} = \mathbf{0}$
 $\langle proof \rangle$

lemma (*in ring*) *one-zero*:
shows (*carrier* $R = \{\mathbf{0}\}$) = ($\mathbf{1} = \mathbf{0}$)
 $\langle proof \rangle$

lemma (*in ring*) *one-not-zero*:
shows (*carrier* $R \neq \{\mathbf{0}\}$) = ($\mathbf{1} \neq \mathbf{0}$)
 $\langle proof \rangle$

Two examples for use of method algebra

lemma
includes *ring* $R +$ *cring* S
shows $\llbracket a \in \text{carrier } R; b \in \text{carrier } R; c \in \text{carrier } S; d \in \text{carrier } S \rrbracket \implies$
 $a \oplus \ominus (a \oplus \ominus b) = b \ \& \ c \otimes_S d = d \otimes_S c$
 $\langle proof \rangle$

lemma
includes *cring*
shows $\llbracket a \in \text{carrier } R; b \in \text{carrier } R \rrbracket \implies a \ominus (a \ominus b) = b$
 $\langle proof \rangle$

9.2.2 Sums over Finite Sets

lemma (*in cring*) *finsum-ldistr*:
 $\llbracket \text{finite } A; a \in \text{carrier } R; f \in A \rightarrow \text{carrier } R \rrbracket \implies$
 $\text{finsum } R \ f \ A \ \otimes \ a = \text{finsum } R \ (\%i. f \ i \ \otimes \ a) \ A$
 $\langle proof \rangle$

lemma (*in cring*) *finsum-rdistr*:
 $\llbracket \text{finite } A; a \in \text{carrier } R; f \in A \rightarrow \text{carrier } R \rrbracket \implies$
 $a \ \otimes \ \text{finsum } R \ f \ A = \text{finsum } R \ (\%i. a \ \otimes \ f \ i) \ A$
 $\langle proof \rangle$

9.3 Integral Domains

lemma (*in domain*) *zero-not-one* [*simp*]:
 $\mathbf{0} \sim = \mathbf{1}$
 $\langle proof \rangle$

lemma (*in domain*) *integral-iff*:
 $\llbracket a \in \text{carrier } R; b \in \text{carrier } R \rrbracket \implies (a \otimes b = \mathbf{0}) = (a = \mathbf{0} \mid b = \mathbf{0})$
 $\langle proof \rangle$

lemma (*in domain*) *m-lcancel*:
assumes *prem*: $a \sim = \mathbf{0}$
and R : $a \in \text{carrier } R \ b \in \text{carrier } R \ c \in \text{carrier } R$
shows $(a \otimes b = a \otimes c) = (b = c)$

<proof>

lemma (in domain) *m-rcancel*:

assumes *prem*: $a \sim = \mathbf{0}$

and *R*: $a \in \text{carrier } R \ b \in \text{carrier } R \ c \in \text{carrier } R$

shows *conc*: $(b \otimes a = c \otimes a) = (b = c)$

<proof>

9.4 Fields

Field would not need to be derived from domain, the properties for domain follow from the assumptions of field

lemma (in cring) *cring-fieldI*:

assumes *field-Units*: $\text{Units } R = \text{carrier } R - \{\mathbf{0}\}$

shows *field* *R*

<proof>

Another variant to show that something is a field

lemma (in cring) *cring-fieldI2*:

assumes *notzero*: $\mathbf{0} \neq \mathbf{1}$

and *inver*: $\bigwedge a. \llbracket a \in \text{carrier } R; a \neq \mathbf{0} \rrbracket \implies \exists b \in \text{carrier } R. a \otimes b = \mathbf{1}$

shows *field* *R*

<proof>

9.5 Morphisms

constdefs (structure *R S*)

ring-hom :: $[(\ 'a, 'm) \text{ ring-scheme}, (\ 'b, 'n) \text{ ring-scheme}] \implies (\ 'a \implies 'b) \text{ set}$

ring-hom *R S* == $\{h. h \in \text{carrier } R \rightarrow \text{carrier } S \ \&$

$(\text{ALL } x \ y. x \in \text{carrier } R \ \& \ y \in \text{carrier } R \ \longrightarrow$

$h(x \otimes y) = h x \otimes_S h y \ \& \ h(x \oplus y) = h x \oplus_S h y) \ \&$

$h \ \mathbf{1} = \mathbf{1}_S\}$

lemma *ring-hom-memI*:

fixes *R* (structure) **and** *S* (structure)

assumes *hom-closed*: $\llbracket x. x \in \text{carrier } R \rrbracket \implies h x \in \text{carrier } S$

and *hom-mult*: $\llbracket x \ y. \llbracket x \in \text{carrier } R; y \in \text{carrier } R \rrbracket \implies$

$h(x \otimes y) = h x \otimes_S h y$

and *hom-add*: $\llbracket x \ y. \llbracket x \in \text{carrier } R; y \in \text{carrier } R \rrbracket \implies$

$h(x \oplus y) = h x \oplus_S h y$

and *hom-one*: $h \ \mathbf{1} = \mathbf{1}_S$

shows $h \in \text{ring-hom } R \ S$

<proof>

lemma *ring-hom-closed*:

$\llbracket h \in \text{ring-hom } R \ S; x \in \text{carrier } R \rrbracket \implies h x \in \text{carrier } S$

<proof>

lemma *ring-hom-mult*:

fixes R (**structure**) **and** S (**structure**)
shows
 $\llbracket h \in \text{ring-hom } R \ S; x \in \text{carrier } R; y \in \text{carrier } R \rrbracket \implies$
 $h (x \otimes y) = h x \otimes_S h y$
 $\langle \text{proof} \rangle$

lemma *ring-hom-add*:
fixes R (**structure**) **and** S (**structure**)
shows
 $\llbracket h \in \text{ring-hom } R \ S; x \in \text{carrier } R; y \in \text{carrier } R \rrbracket \implies$
 $h (x \oplus y) = h x \oplus_S h y$
 $\langle \text{proof} \rangle$

lemma *ring-hom-one*:
fixes R (**structure**) **and** S (**structure**)
shows $h \in \text{ring-hom } R \ S \implies h \mathbf{1} = \mathbf{1}_S$
 $\langle \text{proof} \rangle$

locale *ring-hom-cring* = *cring* R + *cring* S +
fixes h
assumes *homh* [*simp*, *intro*]: $h \in \text{ring-hom } R \ S$
notes *hom-closed* [*simp*, *intro*] = *ring-hom-closed* [*OF homh*]
and *hom-mult* [*simp*] = *ring-hom-mult* [*OF homh*]
and *hom-add* [*simp*] = *ring-hom-add* [*OF homh*]
and *hom-one* [*simp*] = *ring-hom-one* [*OF homh*]

lemma (**in** *ring-hom-cring*) *hom-zero* [*simp*]:
 $h \mathbf{0} = \mathbf{0}_S$
 $\langle \text{proof} \rangle$

lemma (**in** *ring-hom-cring*) *hom-a-inv* [*simp*]:
 $x \in \text{carrier } R \implies h (\ominus x) = \ominus_S h x$
 $\langle \text{proof} \rangle$

lemma (**in** *ring-hom-cring*) *hom-finsum* [*simp*]:
 $\llbracket \text{finite } A; f \in A \rightarrow \text{carrier } R \rrbracket \implies$
 $h (\text{finsum } R \ f \ A) = \text{finsum } S \ (h \circ f) \ A$
 $\langle \text{proof} \rangle$

lemma (**in** *ring-hom-cring*) *hom-finprod*:
 $\llbracket \text{finite } A; f \in A \rightarrow \text{carrier } R \rrbracket \implies$
 $h (\text{finprod } R \ f \ A) = \text{finprod } S \ (h \circ f) \ A$
 $\langle \text{proof} \rangle$

declare *ring-hom-cring.hom-finprod* [*simp*]

lemma *id-ring-hom* [*simp*]:
 $id \in \text{ring-hom } R \ R$
 $\langle \text{proof} \rangle$

end

theory *Module* imports *Ring* begin

10 Modules over an Abelian Group

10.1 Definitions

record ('a, 'b) *module* = 'b *ring* +
smult :: ['a, 'b] => 'b (**infixl** \odot_M 70)

locale *module* = *cring* *R* + *abelian-group* *M* +
assumes *smult-closed* [*simp*, *intro*]:
 [| *a* ∈ *carrier R*; *x* ∈ *carrier M* |] ==> *a* \odot_M *x* ∈ *carrier M*
and *smult-l-distr*:
 [| *a* ∈ *carrier R*; *b* ∈ *carrier R*; *x* ∈ *carrier M* |] ==>
 (*a* \oplus *b*) \odot_M *x* = *a* \odot_M *x* \oplus_M *b* \odot_M *x*
and *smult-r-distr*:
 [| *a* ∈ *carrier R*; *x* ∈ *carrier M*; *y* ∈ *carrier M* |] ==>
a \odot_M (*x* \oplus_M *y*) = *a* \odot_M *x* \oplus_M *a* \odot_M *y*
and *smult-assoc1*:
 [| *a* ∈ *carrier R*; *b* ∈ *carrier R*; *x* ∈ *carrier M* |] ==>
 (*a* \otimes *b*) \odot_M *x* = *a* \odot_M (*b* \odot_M *x*)
and *smult-one* [*simp*]:
x ∈ *carrier M* ==> **1** \odot_M *x* = *x*

locale *algebra* = *module* *R M* + *cring* *M* +
assumes *smult-assoc2*:
 [| *a* ∈ *carrier R*; *x* ∈ *carrier M*; *y* ∈ *carrier M* |] ==>
 (*a* \odot_M *x*) \otimes_M *y* = *a* \odot_M (*x* \otimes_M *y*)

lemma *moduleI*:

fixes *R* (**structure**) **and** *M* (**structure**)
assumes *cring*: *cring* *R*
and *abelian-group*: *abelian-group* *M*
and *smult-closed*:
 !!*a x*. [| *a* ∈ *carrier R*; *x* ∈ *carrier M* |] ==> *a* \odot_M *x* ∈ *carrier M*
and *smult-l-distr*:
 !!*a b x*. [| *a* ∈ *carrier R*; *b* ∈ *carrier R*; *x* ∈ *carrier M* |] ==>
 (*a* \oplus *b*) \odot_M *x* = (*a* \odot_M *x*) \oplus_M (*b* \odot_M *x*)
and *smult-r-distr*:
 !!*a x y*. [| *a* ∈ *carrier R*; *x* ∈ *carrier M*; *y* ∈ *carrier M* |] ==>
a \odot_M (*x* \oplus_M *y*) = (*a* \odot_M *x*) \oplus_M (*a* \odot_M *y*)
and *smult-assoc1*:
 !!*a b x*. [| *a* ∈ *carrier R*; *b* ∈ *carrier R*; *x* ∈ *carrier M* |] ==>
 (*a* \otimes *b*) \odot_M *x* = *a* \odot_M (*b* \odot_M *x*)

and smult-one:
 $!!x. x \in \text{carrier } M \implies \mathbf{1} \odot_M x = x$
shows module R M
 $\langle \text{proof} \rangle$

lemma algebraI:
fixes R (structure) and M (structure)
assumes R -cring: cring R
and M -cring: cring M
and smult-closed:
 $!!a x. [! a \in \text{carrier } R; x \in \text{carrier } M] \implies a \odot_M x \in \text{carrier } M$
and smult-l-distr:
 $!!a b x. [! a \in \text{carrier } R; b \in \text{carrier } R; x \in \text{carrier } M] \implies$
 $(a \oplus b) \odot_M x = (a \odot_M x) \oplus_M (b \odot_M x)$
and smult-r-distr:
 $!!a x y. [! a \in \text{carrier } R; x \in \text{carrier } M; y \in \text{carrier } M] \implies$
 $a \odot_M (x \oplus_M y) = (a \odot_M x) \oplus_M (a \odot_M y)$
and smult-assoc1:
 $!!a b x. [! a \in \text{carrier } R; b \in \text{carrier } R; x \in \text{carrier } M] \implies$
 $(a \otimes b) \odot_M x = a \odot_M (b \odot_M x)$
and smult-one:
 $!!x. x \in \text{carrier } M \implies (\text{one } R) \odot_M x = x$
and smult-assoc2:
 $!!a x y. [! a \in \text{carrier } R; x \in \text{carrier } M; y \in \text{carrier } M] \implies$
 $(a \odot_M x) \otimes_M y = a \odot_M (x \otimes_M y)$
shows algebra R M
 $\langle \text{proof} \rangle$

lemma (in algebra) R -cring:
 $\text{cring } R$
 $\langle \text{proof} \rangle$

lemma (in algebra) M -cring:
 $\text{cring } M$
 $\langle \text{proof} \rangle$

lemma (in algebra) module:
 $\text{module } R$ M
 $\langle \text{proof} \rangle$

10.2 Basic Properties of Algebras

lemma (in algebra) smult-l-null [simp]:
 $x \in \text{carrier } M \implies \mathbf{0} \odot_M x = \mathbf{0}_M$
 $\langle \text{proof} \rangle$

lemma (in algebra) smult-r-null [simp]:
 $a \in \text{carrier } R \implies a \odot_M \mathbf{0}_M = \mathbf{0}_M$
 $\langle \text{proof} \rangle$

lemma (in algebra) *smult-l-minus*:

$[[a \in \text{carrier } R; x \in \text{carrier } M]] \implies (\ominus a) \odot_M x = \ominus_M (a \odot_M x)$
 <proof>

lemma (in algebra) *smult-r-minus*:

$[[a \in \text{carrier } R; x \in \text{carrier } M]] \implies a \odot_M (\ominus_M x) = \ominus_M (a \odot_M x)$
 <proof>

end

theory *UnivPoly* imports *Module* begin

11 Univariate Polynomials

Polynomials are formalised as modules with additional operations for extracting coefficients from polynomials and for obtaining monomials from coefficients and exponents (record *up-ring*). The carrier set is a set of bounded functions from Nat to the coefficient domain. Bounded means that these functions return zero above a certain bound (the degree). There is a chapter on the formalisation of polynomials in the PhD thesis [1], which was implemented with axiomatic type classes. This was later ported to Locales.

11.1 The Constructor for Univariate Polynomials

Functions with finite support.

locale *bound* =

fixes $z :: 'a$
and $n :: \text{nat}$
and $f :: \text{nat} \Rightarrow 'a$
assumes *bound*: $!!m. n < m \implies f m = z$

declare *bound.intro* [*intro!*]

and *bound.bound* [*dest*]

lemma *bound-below*:

assumes *bound*: $\text{bound } z \ m \ f$ **and** *nonzero*: $f \ n \neq z$ **shows** $n \leq m$
 <proof>

record $('a, 'p)$ *up-ring* = $('a, 'p)$ *module* +

monom :: $['a, \text{nat}] \Rightarrow 'p$

coeff :: $['p, \text{nat}] \Rightarrow 'a$

constdefs (structure *R*)

up :: $('a, 'm)$ *ring-scheme* $\Rightarrow (\text{nat} \Rightarrow 'a)$ *set*

```

up R == {f. f ∈ UNIV -> carrier R & (EX n. bound 0 n f)}
UP :: ('a, 'm) ring-scheme => ('a, nat => 'a) up-ring
UP R == (|
  carrier = up R,
  mult = (%p:up R. %q:up R. %n. ⊕ i ∈ {...n}. p i ⊗ q (n-i)),
  one = (%i. if i=0 then 1 else 0),
  zero = (%i. 0),
  add = (%p:up R. %q:up R. %i. p i ⊕ q i),
  smult = (%a:carrier R. %p:up R. %i. a ⊗ p i),
  monom = (%a:carrier R. %n i. if i=n then a else 0),
  coeff = (%p:up R. %n. p n) |)

```

Properties of the set of polynomials *up*.

lemma *mem-upI* [*intro*]:

```

[| !!n. f n ∈ carrier R; EX n. bound (zero R) n f |] ==> f ∈ up R
⟨proof⟩

```

lemma *mem-upD* [*dest*]:

```

f ∈ up R ==> f n ∈ carrier R
⟨proof⟩

```

lemma (*in cring*) *bound-upD* [*dest*]:

```

f ∈ up R ==> EX n. bound 0 n f
⟨proof⟩

```

lemma (*in cring*) *up-one-closed*:

```

(%n. if n = 0 then 1 else 0) ∈ up R
⟨proof⟩

```

lemma (*in cring*) *up-smult-closed*:

```

[| a ∈ carrier R; p ∈ up R |] ==> (%i. a ⊗ p i) ∈ up R
⟨proof⟩

```

lemma (*in cring*) *up-add-closed*:

```

[| p ∈ up R; q ∈ up R |] ==> (%i. p i ⊕ q i) ∈ up R
⟨proof⟩

```

lemma (*in cring*) *up-a-inv-closed*:

```

p ∈ up R ==> (%i. ⊖ (p i)) ∈ up R
⟨proof⟩

```

lemma (*in cring*) *up-mult-closed*:

```

[| p ∈ up R; q ∈ up R |] ==>
(%n. ⊕ i ∈ {...n}. p i ⊗ q (n-i)) ∈ up R
⟨proof⟩

```

11.2 Effect of Operations on Coefficients

locale *UP* =

fixes R (structure) and P (structure)
defines P -def: $P == UP R$

locale UP -cring = $UP + cring R$

locale UP -domain = UP -cring + domain R

Temporarily declare $P \equiv UP R$ as simp rule.

declare (in UP) P -def [simp]

lemma (in UP -cring) *coeff-monom* [simp]:
 $a \in carrier R ==>$
 $coeff P (monom P a m) n = (if m=n then a else 0)$
 <proof>

lemma (in UP -cring) *coeff-zero* [simp]:
 $coeff P 0_P n = 0$
 <proof>

lemma (in UP -cring) *coeff-one* [simp]:
 $coeff P 1_P n = (if n=0 then 1 else 0)$
 <proof>

lemma (in UP -cring) *coeff-smult* [simp]:
 $[| a \in carrier R; p \in carrier P |] ==>$
 $coeff P (a \odot_P p) n = a \otimes coeff P p n$
 <proof>

lemma (in UP -cring) *coeff-add* [simp]:
 $[| p \in carrier P; q \in carrier P |] ==>$
 $coeff P (p \oplus_P q) n = coeff P p n \oplus coeff P q n$
 <proof>

lemma (in UP -cring) *coeff-mult* [simp]:
 $[| p \in carrier P; q \in carrier P |] ==>$
 $coeff P (p \otimes_P q) n = (\bigoplus_{i \in \{..n\}} coeff P p i \otimes coeff P q (n-i))$
 <proof>

lemma (in UP) *up-eqI*:
assumes *prem*: $!!n. coeff P p n = coeff P q n$
and R : $p \in carrier P \ q \in carrier P$
shows $p = q$
 <proof>

11.3 Polynomials Form a Commutative Ring.

Operations are closed over P .

lemma (in UP -cring) *UP-mult-closed* [simp]:
 $[| p \in carrier P; q \in carrier P |] ==> p \otimes_P q \in carrier P$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-one-closed* [simp]:

$\mathbf{1}_P \in carrier\ P$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-zero-closed* [intro, simp]:

$\mathbf{0}_P \in carrier\ P$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-a-closed* [intro, simp]:

$[[\ p \in carrier\ P; q \in carrier\ P \]] \implies p \oplus_P q \in carrier\ P$

$\langle proof \rangle$

lemma (in *UP-cring*) *monom-closed* [simp]:

$a \in carrier\ R \implies monom\ P\ a\ n \in carrier\ P$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-smult-closed* [simp]:

$[[\ a \in carrier\ R; p \in carrier\ P \]] \implies a \odot_P p \in carrier\ P$

$\langle proof \rangle$

lemma (in *UP*) *coeff-closed* [simp]:

$p \in carrier\ P \implies coeff\ P\ p\ n \in carrier\ R$

$\langle proof \rangle$

declare (in *UP*) *P-def* [simp del]

Algebraic ring properties

lemma (in *UP-cring*) *UP-a-assoc*:

assumes $R: p \in carrier\ P\ q \in carrier\ P\ r \in carrier\ P$

shows $(p \oplus_P q) \oplus_P r = p \oplus_P (q \oplus_P r)$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-l-zero* [simp]:

assumes $R: p \in carrier\ P$

shows $\mathbf{0}_P \oplus_P p = p$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-l-neg-ex*:

assumes $R: p \in carrier\ P$

shows $EX\ q : carrier\ P. q \oplus_P p = \mathbf{0}_P$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-a-comm*:

assumes $R: p \in carrier\ P\ q \in carrier\ P$

shows $p \oplus_P q = q \oplus_P p$

$\langle proof \rangle$

lemma (in *UP-cring*) *UP-m-assoc*:
 assumes $R: p \in \text{carrier } P \ q \in \text{carrier } P \ r \in \text{carrier } P$
 shows $(p \otimes_P q) \otimes_P r = p \otimes_P (q \otimes_P r)$
 <proof>

lemma (in *UP-cring*) *UP-l-one* [*simp*]:
 assumes $R: p \in \text{carrier } P$
 shows $\mathbf{1}_P \otimes_P p = p$
 <proof>

lemma (in *UP-cring*) *UP-l-distr*:
 assumes $R: p \in \text{carrier } P \ q \in \text{carrier } P \ r \in \text{carrier } P$
 shows $(p \oplus_P q) \otimes_P r = (p \otimes_P r) \oplus_P (q \otimes_P r)$
 <proof>

lemma (in *UP-cring*) *UP-m-comm*:
 assumes $R: p \in \text{carrier } P \ q \in \text{carrier } P$
 shows $p \otimes_P q = q \otimes_P p$
 <proof>

theorem (in *UP-cring*) *UP-cring*:
 cring P
 <proof>

lemma (in *UP-cring*) *UP-ring*:
 ring P
 <proof>

lemma (in *UP-cring*) *UP-a-inv-closed* [*intro, simp*]:
 $p \in \text{carrier } P \implies \ominus_P p \in \text{carrier } P$
 <proof>

lemma (in *UP-cring*) *coeff-a-inv* [*simp*]:
 assumes $R: p \in \text{carrier } P$
 shows $\text{coeff } P (\ominus_P p) \ n = \ominus (\text{coeff } P \ p \ n)$
 <proof>

Interpretation of lemmas from *cring*. Saves lifting 43 lemmas manually.

interpretation *UP-cring* < *cring* P
 <proof>

11.4 Polynomials Form an Algebra

lemma (in *UP-cring*) *UP-smult-l-distr*:
 $\llbracket a \in \text{carrier } R; b \in \text{carrier } R; p \in \text{carrier } P \rrbracket \implies$
 $(a \oplus b) \odot_P p = a \odot_P p \oplus_P b \odot_P p$
 <proof>

lemma (in *UP-cring*) *UP-smult-r-distr*:
 $\llbracket a \in \text{carrier } R; p \in \text{carrier } P; q \in \text{carrier } P \rrbracket \implies$
 $a \odot_P (p \oplus_P q) = a \odot_P p \oplus_P a \odot_P q$
 ⟨proof⟩

lemma (in *UP-cring*) *UP-smult-assoc1*:
 $\llbracket a \in \text{carrier } R; b \in \text{carrier } R; p \in \text{carrier } P \rrbracket \implies$
 $(a \otimes b) \odot_P p = a \odot_P (b \odot_P p)$
 ⟨proof⟩

lemma (in *UP-cring*) *UP-smult-one* [simp]:
 $p \in \text{carrier } P \implies \mathbf{1} \odot_P p = p$
 ⟨proof⟩

lemma (in *UP-cring*) *UP-smult-assoc2*:
 $\llbracket a \in \text{carrier } R; p \in \text{carrier } P; q \in \text{carrier } P \rrbracket \implies$
 $(a \odot_P p) \otimes_P q = a \odot_P (p \otimes_P q)$
 ⟨proof⟩

Interpretation of lemmas from *algebra*.

lemma (in *cring*) *cring*:
 $\text{cring } R$
 ⟨proof⟩

lemma (in *UP-cring*) *UP-algebra*:
 $\text{algebra } R P$
 ⟨proof⟩

interpretation *UP-cring < algebra R P*
 ⟨proof⟩

11.5 Further Lemmas Involving Monomials

lemma (in *UP-cring*) *monom-zero* [simp]:
 $\text{monom } P \mathbf{0} n = \mathbf{0}_P$
 ⟨proof⟩

lemma (in *UP-cring*) *monom-mult-is-smult*:
assumes $R: a \in \text{carrier } R p \in \text{carrier } P$
shows $\text{monom } P a 0 \otimes_P p = a \odot_P p$
 ⟨proof⟩

lemma (in *UP-cring*) *monom-add* [simp]:
 $\llbracket a \in \text{carrier } R; b \in \text{carrier } R \rrbracket \implies$
 $\text{monom } P (a \oplus b) n = \text{monom } P a n \oplus_P \text{monom } P b n$
 ⟨proof⟩

lemma (in *UP-cring*) *monom-one-Suc*:
 $\text{monom } P \mathbf{1} (\text{Suc } n) = \text{monom } P \mathbf{1} n \otimes_P \text{monom } P \mathbf{1} 1$

<proof>

lemma (in *UP-crng*) *monom-mult-smult*:

$[[a \in \text{carrier } R; b \in \text{carrier } R]] \implies \text{monom } P (a \otimes b) n = a \odot_P \text{monom } P b n$

<proof>

lemma (in *UP-crng*) *monom-one* [*simp*]:

$\text{monom } P \mathbf{1} 0 = \mathbf{1}_P$

<proof>

lemma (in *UP-crng*) *monom-one-mult*:

$\text{monom } P \mathbf{1} (n + m) = \text{monom } P \mathbf{1} n \otimes_P \text{monom } P \mathbf{1} m$

<proof>

lemma (in *UP-crng*) *monom-mult* [*simp*]:

assumes *R*: $a \in \text{carrier } R$ $b \in \text{carrier } R$

shows $\text{monom } P (a \otimes b) (n + m) = \text{monom } P a n \otimes_P \text{monom } P b m$

<proof>

lemma (in *UP-crng*) *monom-a-inv* [*simp*]:

$a \in \text{carrier } R \implies \text{monom } P (\ominus a) n = \ominus_P \text{monom } P a n$

<proof>

lemma (in *UP-crng*) *monom-inj*:

inj-on ($\%a. \text{monom } P a n$) (*carrier* *R*)

<proof>

11.6 The Degree Function

constdefs (structure *R*)

deg :: [$'a, 'm$] *ring-scheme*, $\text{nat} \implies 'a] \implies \text{nat}$

$\text{deg } R p == \text{LEAST } n. \text{bound } \mathbf{0} n (\text{coeff } (UP \ R) p)$

lemma (in *UP-crng*) *deg-aboveI*:

$[[(!m. n < m \implies \text{coeff } P p m = \mathbf{0}); p \in \text{carrier } P]] \implies \text{deg } R p \leq n$

<proof>

lemma (in *UP-crng*) *deg-aboveD*:

assumes $\text{deg } R p < m$ **and** $p \in \text{carrier } P$

shows $\text{coeff } P p m = \mathbf{0}$

<proof>

lemma (in *UP-crng*) *deg-belowI*:

assumes *non-zero*: $n \sim 0 \implies \text{coeff } P p n \sim \mathbf{0}$

and *R*: $p \in \text{carrier } P$

shows $n \leq \text{deg } R p$

— Logically, this is a slightly stronger version of *deg-aboveD*
 ⟨proof⟩

lemma (in *UP-crimg*) *lcoeff-nonzero-deg*:

assumes *deg*: $\text{deg } R \ p \ \sim = 0$ **and** *R*: $p \in \text{carrier } P$

shows $\text{coeff } P \ p \ (\text{deg } R \ p) \ \sim = \mathbf{0}$

⟨proof⟩

lemma (in *UP-crimg*) *lcoeff-nonzero-nonzero*:

assumes *deg*: $\text{deg } R \ p = 0$ **and** *nonzero*: $p \ \sim = \mathbf{0}_P$ **and** *R*: $p \in \text{carrier } P$

shows $\text{coeff } P \ p \ 0 \ \sim = \mathbf{0}$

⟨proof⟩

lemma (in *UP-crimg*) *lcoeff-nonzero*:

assumes *neg*: $p \ \sim = \mathbf{0}_P$ **and** *R*: $p \in \text{carrier } P$

shows $\text{coeff } P \ p \ (\text{deg } R \ p) \ \sim = \mathbf{0}$

⟨proof⟩

lemma (in *UP-crimg*) *deg-eqI*:

[| *m*. $n < m \implies \text{coeff } P \ p \ m = \mathbf{0}$;

!!*n*. $n \ \sim = 0 \implies \text{coeff } P \ p \ n \ \sim = \mathbf{0}$; $p \in \text{carrier } P$ |] $\implies \text{deg } R \ p = n$

⟨proof⟩

Degree and polynomial operations

lemma (in *UP-crimg*) *deg-add* [*simp*]:

assumes *R*: $p \in \text{carrier } P \ q \in \text{carrier } P$

shows $\text{deg } R \ (p \oplus_P q) \leq \max (\text{deg } R \ p) (\text{deg } R \ q)$

⟨proof⟩

lemma (in *UP-crimg*) *deg-monom-le*:

$a \in \text{carrier } R \implies \text{deg } R \ (\text{monom } P \ a \ n) \leq n$

⟨proof⟩

lemma (in *UP-crimg*) *deg-monom* [*simp*]:

[| $a \ \sim = \mathbf{0}$; $a \in \text{carrier } R$ |] $\implies \text{deg } R \ (\text{monom } P \ a \ n) = n$

⟨proof⟩

lemma (in *UP-crimg*) *deg-const* [*simp*]:

assumes *R*: $a \in \text{carrier } R$ **shows** $\text{deg } R \ (\text{monom } P \ a \ 0) = 0$

⟨proof⟩

lemma (in *UP-crimg*) *deg-zero* [*simp*]:

$\text{deg } R \ \mathbf{0}_P = 0$

⟨proof⟩

lemma (in *UP-crimg*) *deg-one* [*simp*]:

$\text{deg } R \ \mathbf{1}_P = 0$

⟨proof⟩

lemma (in *UP-cring*) *deg-uminus* [simp]:
assumes $R: p \in \text{carrier } P$ **shows** $\text{deg } R (\ominus_P p) = \text{deg } R p$
 ⟨proof⟩

lemma (in *UP-domain*) *deg-smult-ring*:
 [$a \in \text{carrier } R; p \in \text{carrier } P$] ==>
 $\text{deg } R (a \odot_P p) \leq (\text{if } a = \mathbf{0} \text{ then } 0 \text{ else } \text{deg } R p)$
 ⟨proof⟩

lemma (in *UP-domain*) *deg-smult* [simp]:
assumes $R: a \in \text{carrier } R; p \in \text{carrier } P$
shows $\text{deg } R (a \odot_P p) = (\text{if } a = \mathbf{0} \text{ then } 0 \text{ else } \text{deg } R p)$
 ⟨proof⟩

lemma (in *UP-cring*) *deg-mult-cring*:
assumes $R: p \in \text{carrier } P; q \in \text{carrier } P$
shows $\text{deg } R (p \otimes_P q) \leq \text{deg } R p + \text{deg } R q$
 ⟨proof⟩

lemma (in *UP-domain*) *deg-mult* [simp]:
 [$p \sim \mathbf{0}_P; q \sim \mathbf{0}_P; p \in \text{carrier } P; q \in \text{carrier } P$] ==>
 $\text{deg } R (p \otimes_P q) = \text{deg } R p + \text{deg } R q$
 ⟨proof⟩

lemma (in *UP-cring*) *coeff-finsum*:
assumes *fin*: *finite* A
shows $p \in A \rightarrow \text{carrier } P ==>$
 $\text{coeff } P (\text{finsum } P p A) k = (\bigoplus i \in A. \text{coeff } P (p i) k)$
 ⟨proof⟩

lemma (in *UP-cring*) *up-repr*:
assumes $R: p \in \text{carrier } P$
shows $(\bigoplus_P i \in \{\dots, \text{deg } R p\}. \text{monom } P (\text{coeff } P p i) i) = p$
 ⟨proof⟩

lemma (in *UP-cring*) *up-repr-le*:
 [$\text{deg } R p \leq n; p \in \text{carrier } P$] ==>
 $(\bigoplus_P i \in \{\dots, n\}. \text{monom } P (\text{coeff } P p i) i) = p$
 ⟨proof⟩

11.7 Polynomials over Integral Domains

lemma *domainI*:
assumes *cring*: *cring* R
and *one-not-zero*: $\text{one } R \sim \text{zero } R$
and *integral*: $\forall a b. [\text{mult } R a b = \text{zero } R; a \in \text{carrier } R;$
 $b \in \text{carrier } R] ==> a = \text{zero } R \mid b = \text{zero } R$
shows *domain* R
 ⟨proof⟩

lemma (in *UP-domain*) *UP-one-not-zero*:

$\mathbf{1}_P \sim = \mathbf{0}_P$
 ⟨proof⟩

lemma (in *UP-domain*) *UP-integral*:

$\llbracket p \otimes_P q = \mathbf{0}_P; p \in \text{carrier } P; q \in \text{carrier } P \rrbracket \implies p = \mathbf{0}_P \mid q = \mathbf{0}_P$
 ⟨proof⟩

theorem (in *UP-domain*) *UP-domain*:

domain P
 ⟨proof⟩

Interpretation of theorems from *domain*.

interpretation *UP-domain < domain P*

⟨proof⟩

11.8 The Evaluation Homomorphism and Universal Property

theorem (in *cring*) *diagonal-sum*:

$\llbracket f \in \{..n + m::\text{nat}\} \rightarrow \text{carrier } R; g \in \{..n + m\} \rightarrow \text{carrier } R \rrbracket \implies$
 $(\bigoplus k \in \{..n + m\}. \bigoplus i \in \{..k\}. f i \otimes g (k - i)) =$
 $(\bigoplus k \in \{..n + m\}. \bigoplus i \in \{..n + m - k\}. f k \otimes g i)$
 ⟨proof⟩

lemma (in *abelian-monoid*) *boundD-carrier*:

$\llbracket \text{bound } \mathbf{0} \ n \ f; n < m \rrbracket \implies f \ m \in \text{carrier } G$
 ⟨proof⟩

theorem (in *cring*) *cauchy-product*:

assumes *bf: bound 0 n f and bg: bound 0 m g*
and *Rf: f ∈ {..n} → carrier R and Rg: g ∈ {..m} → carrier R*
shows $(\bigoplus k \in \{..n + m\}. \bigoplus i \in \{..k\}. f i \otimes g (k - i)) =$
 $(\bigoplus i \in \{..n\}. f i) \otimes (\bigoplus i \in \{..m\}. g i)$
 ⟨proof⟩

lemma (in *UP-cring*) *const-ring-hom*:

$(\%a. \text{monom } P \ a \ 0) \in \text{ring-hom } R \ P$
 ⟨proof⟩

constdefs (structure *S*)

eval :: $[('a, 'm) \text{ ring-scheme}, ('b, 'n) \text{ ring-scheme},$
 $'a \Rightarrow 'b, 'b, \text{nat} \Rightarrow 'a] \Rightarrow 'b$
eval *R S phi s* == $\lambda p \in \text{carrier } (UP \ R).$
 $\bigoplus i \in \{..deg \ R \ p\}. \text{phi } (\text{coeff } (UP \ R) \ p \ i) \otimes s \ (^) \ i$

lemma (in *UP*) *eval-on-carrier*:

fixes S (**structure**)
shows $p \in \text{carrier } P \implies$
 $\text{eval } R \ S \ \text{phi } s \ p = (\bigoplus_S i \in \{.. \text{deg } R \ p\}. \text{phi } (\text{coeff } P \ p \ i) \otimes_S s \ (\cdot)_S i)$
 $\langle \text{proof} \rangle$

lemma (**in** UP) *eval-extensional*:
 $\text{eval } R \ S \ \text{phi } p \in \text{extensional } (\text{carrier } P)$
 $\langle \text{proof} \rangle$

The universal property of the polynomial ring

locale $UP\text{-pre-univ-prop} = \text{ring-hom-crimg } R \ S \ h + UP\text{-crimg } R \ P$

locale $UP\text{-univ-prop} = UP\text{-pre-univ-prop} +$
fixes s **and** $Eval$
assumes $\text{indet-img-carrier } [\text{simp}, \text{intro}]: s \in \text{carrier } S$
defines $Eval\text{-def}: Eval == \text{eval } R \ S \ h \ s$

theorem (**in** $UP\text{-pre-univ-prop}$) *eval-ring-hom*:
assumes $S: s \in \text{carrier } S$
shows $\text{eval } R \ S \ h \ s \in \text{ring-hom } P \ S$
 $\langle \text{proof} \rangle$

Interpretation of ring homomorphism lemmas.

interpretation $UP\text{-univ-prop} < \text{ring-hom-crimg } P \ S \ Eval$
 $\langle \text{proof} \rangle$

Further properties of the evaluation homomorphism.

The following lemma could be proved in $UP\text{-crimg}$ with the additional assumption that h is closed.

lemma (**in** $UP\text{-pre-univ-prop}$) *eval-const*:
 $[[s \in \text{carrier } S; r \in \text{carrier } R]] \implies \text{eval } R \ S \ h \ s \ (\text{monom } P \ r \ 0) = h \ r$
 $\langle \text{proof} \rangle$

The following proof is complicated by the fact that in arbitrary rings one might have $\mathbf{1}_R = \mathbf{0}_R$.

lemma (**in** $UP\text{-pre-univ-prop}$) *eval-monom1*:
assumes $S: s \in \text{carrier } S$
shows $\text{eval } R \ S \ h \ s \ (\text{monom } P \ \mathbf{1} \ 1) = s$
 $\langle \text{proof} \rangle$

lemma (**in** $UP\text{-crimg}$) *monom-pow*:
assumes $R: a \in \text{carrier } R$
shows $(\text{monom } P \ a \ n) \ (\cdot)_P \ m = \text{monom } P \ (a \ (\cdot)_m) \ (n * m)$
 $\langle \text{proof} \rangle$

lemma (**in** ring-hom-crimg) *hom-pow* [*simp*]:
 $x \in \text{carrier } R \implies h \ (x \ (\cdot)_n) = h \ x \ (\cdot)_S \ (n::\text{nat})$

<proof>

lemma (in *UP-univ-prop*) *Eval-monom*:
 $r \in \text{carrier } R \implies \text{Eval } (\text{monom } P \ r \ n) = h \ r \otimes_S s \ (\wedge)_S \ n$
<proof>

lemma (in *UP-pre-univ-prop*) *eval-monom*:
assumes $R: r \in \text{carrier } R$ **and** $S: s \in \text{carrier } S$
shows $\text{eval } R \ S \ h \ s \ (\text{monom } P \ r \ n) = h \ r \otimes_S s \ (\wedge)_S \ n$
<proof>

lemma (in *UP-univ-prop*) *Eval-smult*:
 $[[r \in \text{carrier } R; p \in \text{carrier } P]] \implies \text{Eval } (r \odot_P p) = h \ r \otimes_S \text{Eval } p$
<proof>

lemma *ring-hom-cringI*:
assumes *cring* R
and *cring* S
and $h \in \text{ring-hom } R \ S$
shows *ring-hom-cring* $R \ S \ h$
<proof>

lemma (in *UP-pre-univ-prop*) *UP-hom-unique*:
includes *ring-hom-cring* $P \ S \ \text{Phi}$
assumes $\text{Phi}: \text{Phi } (\text{monom } P \ \mathbf{1} \ (\text{Suc } 0)) = s$
 $!!r. r \in \text{carrier } R \implies \text{Phi } (\text{monom } P \ r \ 0) = h \ r$
includes *ring-hom-cring* $P \ S \ \text{Psi}$
assumes $\text{Psi}: \text{Psi } (\text{monom } P \ \mathbf{1} \ (\text{Suc } 0)) = s$
 $!!r. r \in \text{carrier } R \implies \text{Psi } (\text{monom } P \ r \ 0) = h \ r$
and $P: p \in \text{carrier } P$ **and** $S: s \in \text{carrier } S$
shows $\text{Phi } p = \text{Psi } p$
<proof>

lemma (in *UP-pre-univ-prop*) *ring-homD*:
assumes $\text{Phi}: \text{Phi} \in \text{ring-hom } P \ S$
shows *ring-hom-cring* $P \ S \ \text{Phi}$
<proof>

theorem (in *UP-pre-univ-prop*) *UP-universal-property*:
assumes $S: s \in \text{carrier } S$
shows $\text{EX! } \text{Phi}. \text{Phi} \in \text{ring-hom } P \ S \cap \text{extensional } (\text{carrier } P) \ \&$
 $\text{Phi } (\text{monom } P \ \mathbf{1} \ 1) = s \ \&$
 $(\text{ALL } r : \text{carrier } R. \text{Phi } (\text{monom } P \ r \ 0) = h \ r)$
<proof>

11.9 Sample Application of Evaluation Homomorphism

lemma *UP-pre-univ-propI*:
assumes *cring* R

```

and cring S
and h  $\in$  ring-hom R S
shows UP-pre-univ-prop R S h
  <proof>

```

constdefs

```

INTEG :: int ring
INTEG == (| carrier = UNIV, mult = op *, one = 1, zero = 0, add = op +
|)

```

lemma *INTEG-cring*:

```

cring INTEG
  <proof>

```

lemma *INTEG-id-eval*:

```

UP-pre-univ-prop INTEG INTEG id
  <proof>

```

Interpretation now enables to import all theorems and lemmas valid in the context of homomorphisms between *INTEG* and *UP INTEG* globally.

interpretation *INTEG*: *UP-pre-univ-prop* [*INTEG INTEG id*]

```

  <proof>

```

lemma *INTEG-closed* [*intro*, *simp*]:

```

z  $\in$  carrier INTEG
  <proof>

```

lemma *INTEG-mult* [*simp*]:

```

mult INTEG z w = z * w
  <proof>

```

lemma *INTEG-pow* [*simp*]:

```

pow INTEG z n = z ^ n
  <proof>

```

lemma *eval INTEG INTEG id 10* (*monom* (*UP INTEG*) 5 2) = 500

```

  <proof>

```

end

theory *AbelCoset*

imports *Coset Ring*

begin

12 More Lifting from Groups to Abelian Groups

12.1 Definitions

Hiding $\langle + \rangle$ from *Sum-Type* until I come up with better syntax here

hide *const Plus*

constdefs (structure *G*)

a-r-coset :: $[-, 'a \text{ set}, 'a] \Rightarrow 'a \text{ set}$ (**infixl** $\langle + \rangle_1$ 60)

a-r-coset *G* \equiv *r-coset* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)

a-l-coset :: $[-, 'a, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (**infixl** $\langle + \rangle_1$ 60)

a-l-coset *G* \equiv *l-coset* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)

A-RCOSETS :: $[-, 'a \text{ set}] \Rightarrow ('a \text{ set}) \text{ set}$ (*a'-rcosets1* - [81] 80)

A-RCOSETS *G H* \equiv *RCOSETS* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor) *H*

set-add :: $[-, 'a \text{ set}, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (**infixl** $\langle + \rangle_1$ 60)

set-add *G* \equiv *set-mult* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)

A-SET-INV :: $[-, 'a \text{ set}] \Rightarrow 'a \text{ set}$ (*a'-set'-inv1* - [81] 80)

A-SET-INV *G H* \equiv *SET-INV* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor) *H*

constdefs (structure *G*)

a-r-congruent :: $[('a, 'b) \text{ ring-scheme}, 'a \text{ set}] \Rightarrow ('a * 'a) \text{ set}$
(*racong1* -)

a-r-congruent *G* \equiv *r-congruent* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)

constdefs

A-FactGroup :: $[('a, 'b) \text{ ring-scheme}, 'a \text{ set}] \Rightarrow ('a \text{ set}) \text{ monoid}$
(**infixl** *A'-Mod* 65)

— Actually defined for groups rather than monoids

A-FactGroup *G H* \equiv *FactGroup* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor) *H*

constdefs

a-kernel :: $('a, 'm) \text{ ring-scheme} \Rightarrow ('b, 'n) \text{ ring-scheme} \Rightarrow$
 $('a \Rightarrow 'b) \Rightarrow 'a \text{ set}$

— the kernel of a homomorphism (additive)

a-kernel *G H h* \equiv *kernel* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)
(\lfloor *carrier* = *carrier H*, *mult* = *add H*, *one* = *zero H* \rfloor) *h*

locale *abelian-group-hom* = *abelian-group G* + *abelian-group H* + *var h* +

assumes *a-group-hom*: *group-hom* (\lfloor *carrier* = *carrier G*, *mult* = *add G*, *one* = *zero G* \rfloor)

(\lfloor *carrier* = *carrier H*, *mult* = *add H*, *one* = *zero H* \rfloor) *h*

lemmas *a-r-coset-defs* =
a-r-coset-def r-coset-def

lemma *a-r-coset-def'*:
includes *struct G*
shows $H +> a \equiv \bigcup h \in H. \{h \oplus a\}$
<proof>

lemmas *a-l-coset-defs* =
a-l-coset-def l-coset-def

lemma *a-l-coset-def'*:
includes *struct G*
shows $a <+ H \equiv \bigcup h \in H. \{a \oplus h\}$
<proof>

lemmas *A-RCOSETS-defs* =
A-RCOSETS-def RCOSETS-def

lemma *A-RCOSETS-def'*:
includes *struct G*
shows $a\text{-rcosets } H \equiv \bigcup a \in \text{carrier } G. \{H +> a\}$
<proof>

lemmas *set-add-defs* =
set-add-def set-mult-def

lemma *set-add-def'*:
includes *struct G*
shows $H <+> K \equiv \bigcup h \in H. \bigcup k \in K. \{h \oplus k\}$
<proof>

lemmas *A-SET-INV-defs* =
A-SET-INV-def SET-INV-def

lemma *A-SET-INV-def'*:
includes *struct G*
shows $a\text{-set-inv } H \equiv \bigcup h \in H. \{\ominus h\}$
<proof>

12.2 Cosets

lemma (in *abelian-group*) *a-coset-add-assoc*:
 $[[M \subseteq \text{carrier } G; g \in \text{carrier } G; h \in \text{carrier } G]]$
 $\implies (M +> g) +> h = M +> (g \oplus h)$
<proof>

lemma (in *abelian-group*) *a-coset-add-zero* [*simp*]:

$M \subseteq \text{carrier } G \implies M +> \mathbf{0} = M$
 ⟨proof⟩

lemma (in *abelian-group*) *a-coset-add-inv1*:

$\llbracket M +> (x \oplus (\ominus y)) = M; x \in \text{carrier } G; y \in \text{carrier } G;$
 $M \subseteq \text{carrier } G \rrbracket \implies M +> x = M +> y$

⟨proof⟩

lemma (in *abelian-group*) *a-coset-add-inv2*:

$\llbracket M +> x = M +> y; x \in \text{carrier } G; y \in \text{carrier } G; M \subseteq \text{carrier } G \rrbracket$
 $\implies M +> (x \oplus (\ominus y)) = M$

⟨proof⟩

lemma (in *abelian-group*) *a-coset-join1*:

$\llbracket H +> x = H; x \in \text{carrier } G; \text{subgroup } H (\text{carrier} = \text{carrier } G, \text{mult} =$
 $\text{add } G, \text{one} = \text{zero } G) \rrbracket \implies x \in H$

⟨proof⟩

lemma (in *abelian-group*) *a-solve-equation*:

$\llbracket \text{subgroup } H (\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G); x \in H; y$
 $\in H \rrbracket \implies \exists h \in H. y = h \oplus x$

⟨proof⟩

lemma (in *abelian-group*) *a-repr-independence*:

$\llbracket y \in H +> x; x \in \text{carrier } G; \text{subgroup } H (\text{carrier} = \text{carrier } G, \text{mult} = \text{add}$
 $G, \text{one} = \text{zero } G) \rrbracket \implies H +> x = H +> y$

⟨proof⟩

lemma (in *abelian-group*) *a-coset-join2*:

$\llbracket x \in \text{carrier } G; \text{subgroup } H (\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero}$
 $G); x \in H \rrbracket \implies H +> x = H$

⟨proof⟩

lemma (in *abelian-monoid*) *a-r-coset-subset-G*:

$\llbracket H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies H +> x \subseteq \text{carrier } G$

⟨proof⟩

lemma (in *abelian-group*) *a-rcosI*:

$\llbracket h \in H; H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies h \oplus x \in H +> x$

⟨proof⟩

lemma (in *abelian-group*) *a-rcosetsI*:

$\llbracket H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies H +> x \in \text{a-rcosets } H$

⟨proof⟩

Really needed?

lemma (in *abelian-group*) *a-transpose-inv*:

$\llbracket x \oplus y = z; x \in \text{carrier } G; y \in \text{carrier } G; z \in \text{carrier } G \rrbracket$
 $\implies (\ominus x) \oplus z = y$

<proof>

12.3 Subgroups

locale *additive-subgroup* = *var* H + *struct* G +
assumes *a-subgroup*: *subgroup* H (\langle *carrier* = *carrier* G , *mult* = *add* G , *one* =
zero G \rangle)

lemma (**in** *additive-subgroup*) *is-additive-subgroup*:

shows *additive-subgroup* H G

<proof>

lemma *additive-subgroupI*:

includes *struct* G

assumes *a-subgroup*: *subgroup* H (\langle *carrier* = *carrier* G , *mult* = *add* G , *one* =
zero G \rangle)

shows *additive-subgroup* H G

<proof>

lemma (**in** *additive-subgroup*) *a-subset*:

$H \subseteq$ *carrier* G

<proof>

lemma (**in** *additive-subgroup*) *a-closed* [*intro*, *simp*]:

$\llbracket x \in H; y \in H \rrbracket \implies x \oplus y \in H$

<proof>

lemma (**in** *additive-subgroup*) *zero-closed* [*simp*]:

$\mathbf{0} \in H$

<proof>

lemma (**in** *additive-subgroup*) *a-inv-closed* [*intro*, *simp*]:

$x \in H \implies \ominus x \in H$

<proof>

12.4 Normal additive subgroups

12.4.1 Definition of *abelian-subgroup*

Every subgroup of an *abelian-group* is normal

locale *abelian-subgroup* = *additive-subgroup* H G + *abelian-group* G +
assumes *a-normal*: *normal* H (\langle *carrier* = *carrier* G , *mult* = *add* G , *one* = *zero*
 G \rangle)

lemma (**in** *abelian-subgroup*) *is-abelian-subgroup*:

shows *abelian-subgroup* H G

<proof>

lemma *abelian-subgroupI*:

assumes *a-normal*: *normal* H ($\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G$)
and *a-comm*: $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket \implies x \oplus_G y = y \oplus_G x$
shows *abelian-subgroup* $H G$
 $\langle \text{proof} \rangle$

lemma *abelian-subgroupI2*:
includes *struct* G
assumes *a-comm-group*: *comm-group* ($\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G$)
and *a-subgroup*: *subgroup* H ($\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G$)
shows *abelian-subgroup* $H G$
 $\langle \text{proof} \rangle$

lemma *abelian-subgroupI3*:
includes *struct* G
assumes *asg*: *additive-subgroup* $H G$
and *ag*: *abelian-group* G
shows *abelian-subgroup* $H G$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-subgroup*) *a-coset-eq*:
 $(\forall x \in \text{carrier } G. H <+ x = x <+ H)$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-subgroup*) *a-inv-op-closed1*:
shows $\llbracket x \in \text{carrier } G; h \in H \rrbracket \implies (\ominus x) \oplus h \oplus x \in H$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-subgroup*) *a-inv-op-closed2*:
shows $\llbracket x \in \text{carrier } G; h \in H \rrbracket \implies x \oplus h \oplus (\ominus x) \in H$
 $\langle \text{proof} \rangle$

Alternative characterization of normal subgroups

lemma (**in** *abelian-group*) *a-normal-inv-iff*:
 $(N \triangleleft (\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G)) =$
 $(\text{subgroup } N (\text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G) \ \& \ (\forall x \in \text{carrier } G. \forall h \in N. x \oplus h \oplus (\ominus x) \in N))$
(is - = ?rhs)
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-group*) *a-lcos-m-assoc*:
 $\llbracket M \subseteq \text{carrier } G; g \in \text{carrier } G; h \in \text{carrier } G \rrbracket$
 $\implies g <+ (h <+ M) = (g \oplus h) <+ M$
 $\langle \text{proof} \rangle$

lemma (**in** *abelian-group*) *a-lcos-mult-one*:

$M \subseteq \text{carrier } G \implies \mathbf{0} <+ M = M$
 ⟨proof⟩

lemma (in *abelian-group*) *a-l-coset-subset-G*:
 $\llbracket H \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket \implies x <+ H \subseteq \text{carrier } G$
 ⟨proof⟩

lemma (in *abelian-group*) *a-l-coset-swap*:
 $\llbracket y \in x <+ H; x \in \text{carrier } G; \text{ subgroup } H \ (\text{carrier} = \text{carrier } G, \text{ mult} = \text{add } G, \text{ one} = \text{zero } G) \rrbracket \implies x \in y <+ H$
 ⟨proof⟩

lemma (in *abelian-group*) *a-l-coset-carrier*:
 $\llbracket y \in x <+ H; x \in \text{carrier } G; \text{ subgroup } H \ (\text{carrier} = \text{carrier } G, \text{ mult} = \text{add } G, \text{ one} = \text{zero } G) \rrbracket \implies y \in \text{carrier } G$
 ⟨proof⟩

lemma (in *abelian-group*) *a-l-repr-imp-subset*:
assumes $y: y \in x <+ H$ **and** $x: x \in \text{carrier } G$ **and** $sb: \text{subgroup } H \ (\text{carrier} = \text{carrier } G, \text{ mult} = \text{add } G, \text{ one} = \text{zero } G)$
shows $y <+ H \subseteq x <+ H$
 ⟨proof⟩

lemma (in *abelian-group*) *a-l-repr-independence*:
assumes $y: y \in x <+ H$ **and** $x: x \in \text{carrier } G$ **and** $sb: \text{subgroup } H \ (\text{carrier} = \text{carrier } G, \text{ mult} = \text{add } G, \text{ one} = \text{zero } G)$
shows $x <+ H = y <+ H$
 ⟨proof⟩

lemma (in *abelian-group*) *setadd-subset-G*:
 $\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G \rrbracket \implies H <+> K \subseteq \text{carrier } G$
 ⟨proof⟩

lemma (in *abelian-group*) *subgroup-add-id*: $\text{subgroup } H \ (\text{carrier} = \text{carrier } G, \text{ mult} = \text{add } G, \text{ one} = \text{zero } G) \implies H <+> H = H$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcos-inv*:
assumes $x: x \in \text{carrier } G$
shows $a\text{-set-inv } (H +> x) = H +> (\ominus x)$
 ⟨proof⟩

lemma (in *abelian-group*) *a-setmult-rcos-assoc*:
 $\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket$
 $\implies H <+> (K +> x) = (H <+> K) +> x$
 ⟨proof⟩

lemma (in *abelian-group*) *a-rcos-assoc-lcos*:
 $\llbracket H \subseteq \text{carrier } G; K \subseteq \text{carrier } G; x \in \text{carrier } G \rrbracket$
 $\implies (H +> x) <+> K = H <+> (x <+ K)$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcos-sum*:
 $\llbracket x \in \text{carrier } G; y \in \text{carrier } G \rrbracket$
 $\implies (H +> x) <+> (H +> y) = H +> (x \oplus y)$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *rcosets-add-eq*:
 $M \in \text{a-rcosets } H \implies H <+> M = M$
 — generalizes *subgroup-mult-id*
 ⟨proof⟩

12.5 Congruence Relation

lemma (in *abelian-subgroup*) *a-equiv-rcong*:
 shows *equiv* (*carrier G*) (*racong H*)
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-l-coset-eq-rcong*:
 assumes $a \in \text{carrier } G$
 shows $a <+ H = \text{racong } H \text{ “ } \{a\}$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcos-equation*:
 shows
 $\llbracket ha \oplus a = h \oplus b; a \in \text{carrier } G; b \in \text{carrier } G;$
 $h \in H; ha \in H; hb \in H \rrbracket$
 $\implies hb \oplus a \in (\bigcup h \in H. \{h \oplus b\})$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcos-disjoint*:
 shows $\llbracket a \in \text{a-rcosets } H; b \in \text{a-rcosets } H; a \neq b \rrbracket \implies a \cap b = \{\}$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcos-self*:
 shows $x \in \text{carrier } G \implies x \in H +> x$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-rcosets-part-G*:
 shows $\bigcup (\text{a-rcosets } H) = \text{carrier } G$
 ⟨proof⟩

lemma (in *abelian-subgroup*) *a-cosets-finite*:
 $\llbracket c \in \text{a-rcosets } H; H \subseteq \text{carrier } G; \text{finite } (\text{carrier } G) \rrbracket \implies \text{finite } c$
 ⟨proof⟩

lemma (in *abelian-group*) *a-card-cosets-equal*:
 $\llbracket c \in a\text{-rcosets } H; H \subseteq \text{carrier } G; \text{finite}(\text{carrier } G) \rrbracket$
 $\implies \text{card } c = \text{card } H$
 ⟨*proof*⟩

lemma (in *abelian-group*) *rcosets-subset-PowG*:
 $\text{additive-subgroup } H \ G \implies a\text{-rcosets } H \subseteq \text{Pow}(\text{carrier } G)$
 ⟨*proof*⟩

theorem (in *abelian-group*) *a-lagrange*:
 $\llbracket \text{finite}(\text{carrier } G); \text{additive-subgroup } H \ G \rrbracket$
 $\implies \text{card}(a\text{-rcosets } H) * \text{card}(H) = \text{order}(G)$
 ⟨*proof*⟩

12.6 Factorization

lemmas *A-FactGroup-defs* = *A-FactGroup-def* *FactGroup-def*

lemma *A-FactGroup-def'*:
includes *struct* *G*
shows $G \ A\text{-Mod } H \equiv (\text{carrier} = a\text{-rcosets}_G \ H, \text{mult} = \text{set-add } G, \text{one} = H)$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-setmult-closed*:
 $\llbracket K1 \in a\text{-rcosets } H; K2 \in a\text{-rcosets } H \rrbracket \implies K1 \langle + \rangle K2 \in a\text{-rcosets } H$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-setinv-closed*:
 $K \in a\text{-rcosets } H \implies a\text{-set-inv } K \in a\text{-rcosets } H$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-rcosets-assoc*:
 $\llbracket M1 \in a\text{-rcosets } H; M2 \in a\text{-rcosets } H; M3 \in a\text{-rcosets } H \rrbracket$
 $\implies M1 \langle + \rangle M2 \langle + \rangle M3 = M1 \langle + \rangle (M2 \langle + \rangle M3)$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-subgroup-in-rcosets*:
 $H \in a\text{-rcosets } H$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-rcosets-inv-mult-group-eq*:
 $M \in a\text{-rcosets } H \implies a\text{-set-inv } M \langle + \rangle M = H$
 ⟨*proof*⟩

theorem (in *abelian-subgroup*) *a-factorgroup-is-group*:
 $\text{group } (G \ A\text{-Mod } H)$
 ⟨*proof*⟩

Since the Factorization is based on an *abelian* subgroup, is results in a

commutative group

theorem (in *abelian-subgroup*) *a-factorgroup-is-comm-group*:
comm-group (G *A-Mod* H)
 ⟨*proof*⟩

lemma *add-A-FactGroup* [*simp*]: $X \otimes_{(G \text{ A-Mod } H)} X' = X <+>_G X'$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-inv-FactGroup*:
 $X \in \text{carrier } (G \text{ A-Mod } H) \implies \text{inv}_{G \text{ A-Mod } H} X = \text{a-set-inv } X$
 ⟨*proof*⟩

The coset map is a homomorphism from G to the quotient group $G \text{ Mod } H$

lemma (in *abelian-subgroup*) *a-r-coset-hom-A-Mod*:
 $(\lambda a. H <+> a) \in \text{hom } (| \text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G |) (G \text{ A-Mod } H)$
 ⟨*proof*⟩

The isomorphism theorems have been omitted from lifting, at least for now

12.7 The First Isomorphism Theorem

The quotient by the kernel of a homomorphism is isomorphic to the range of that homomorphism.

lemmas *a-kernel-defs* =
a-kernel-def *kernel-def*

lemma *a-kernel-def'*:
 $\text{a-kernel } R \ S \ h \equiv \{x \in \text{carrier } R. h \ x = \mathbf{0}_S\}$
 ⟨*proof*⟩

12.8 Homomorphisms

lemma *abelian-group-homI*:
includes *abelian-group* G
includes *abelian-group* H
assumes *a-group-hom*: *group-hom* ($| \text{carrier} = \text{carrier } G, \text{mult} = \text{add } G, \text{one} = \text{zero } G |$)
 $(| \text{carrier} = \text{carrier } H, \text{mult} = \text{add } H, \text{one} = \text{zero } H |) \ h$
shows *abelian-group-hom* $G \ H \ h$
 ⟨*proof*⟩

lemma (in *abelian-group-hom*) *is-abelian-group-hom*:
abelian-group-hom $G \ H \ h$
 ⟨*proof*⟩

lemma (in *abelian-group-hom*) *hom-add* [*simp*]:
 $[| \ x : \text{carrier } G; \ y : \text{carrier } G \ |]$

$\implies h (x \oplus_G y) = h x \oplus_H h y$
 <proof>

lemma (in *abelian-group-hom*) *hom-closed* [simp]:
 $x \in \text{carrier } G \implies h x \in \text{carrier } H$
 <proof>

lemma (in *abelian-group-hom*) *zero-closed* [simp]:
 $h \mathbf{0} \in \text{carrier } H$
 <proof>

lemma (in *abelian-group-hom*) *hom-zero* [simp]:
 $h \mathbf{0} = \mathbf{0}_H$
 <proof>

lemma (in *abelian-group-hom*) *a-inv-closed* [simp]:
 $x \in \text{carrier } G \implies h (\ominus x) \in \text{carrier } H$
 <proof>

lemma (in *abelian-group-hom*) *hom-a-inv* [simp]:
 $x \in \text{carrier } G \implies h (\ominus x) = \ominus_H (h x)$
 <proof>

lemma (in *abelian-group-hom*) *additive-subgroup-a-kernel*:
additive-subgroup (a-kernel G H h) G
 <proof>

The kernel of a homomorphism is an abelian subgroup

lemma (in *abelian-group-hom*) *abelian-subgroup-a-kernel*:
abelian-subgroup (a-kernel G H h) G
 <proof>

lemma (in *abelian-group-hom*) *A-FactGroup-nonempty*:
assumes $X: X \in \text{carrier } (G \text{ A-Mod } a\text{-kernel } G \text{ H } h)$
shows $X \neq \{\}$
 <proof>

lemma (in *abelian-group-hom*) *FactGroup-contents-mem*:
assumes $X: X \in \text{carrier } (G \text{ A-Mod } (a\text{-kernel } G \text{ H } h))$
shows $\text{contents } (h'X) \in \text{carrier } H$
 <proof>

lemma (in *abelian-group-hom*) *A-FactGroup-hom*:
 $(\lambda X. \text{contents } (h'X)) \in \text{hom } (G \text{ A-Mod } (a\text{-kernel } G \text{ H } h))$
 $(\text{carrier} = \text{carrier } H, \text{mult} = \text{add } H, \text{one} = \text{zero } H)$
 <proof>

lemma (in *abelian-group-hom*) *A-FactGroup-inj-on*:
inj-on $(\lambda X. \text{contents } (h'X)) (\text{carrier } (G \text{ A-Mod } a\text{-kernel } G \text{ H } h))$

<proof>

If the homomorphism h is onto H , then so is the homomorphism from the quotient group

lemma (in *abelian-group-hom*) *A-FactGroup-onto*:

assumes $h: h \text{ ' carrier } G = \text{carrier } H$

shows $(\lambda X. \text{contents } (h \text{ ' } X)) \text{ ' carrier } (G \text{ A-Mod } a\text{-kernel } G \text{ H } h) = \text{carrier } H$

<proof>

If h is a homomorphism from G onto H , then the quotient group $G \text{ Mod } \text{kernel } G \text{ H } h$ is isomorphic to H .

theorem (in *abelian-group-hom*) *A-FactGroup-iso*:

$h \text{ ' carrier } G = \text{carrier } H$

$\implies (\lambda X. \text{contents } (h \text{ ' } X)) \in (G \text{ A-Mod } (a\text{-kernel } G \text{ H } h)) \cong$

$(| \text{carrier} = \text{carrier } H, \text{mult} = \text{add } H, \text{one} = \text{zero } H |)$

<proof>

13 Lemmas Lifted from CosetExt.thy

Not everything from `CosetExt.thy` is lifted here.

13.1 General Lemmas from AlgebraExt.thy

lemma (in *additive-subgroup*) *a-Hcarr [simp]*:

assumes $hH: h \in H$

shows $h \in \text{carrier } G$

<proof>

13.2 Lemmas for Right Cosets

lemma (in *abelian-subgroup*) *a-elemrcos-carrier*:

assumes $acarr: a \in \text{carrier } G$

and $a': a' \in H \text{ +> } a$

shows $a' \in \text{carrier } G$

<proof>

lemma (in *abelian-subgroup*) *a-rcos-const*:

assumes $hH: h \in H$

shows $H \text{ +> } h = H$

<proof>

lemma (in *abelian-subgroup*) *a-rcos-module-imp*:

assumes $xcarr: x \in \text{carrier } G$

and $x'cos: x' \in H \text{ +> } x$

shows $(x' \oplus \ominus x) \in H$

<proof>

lemma (in *abelian-subgroup*) *a-rcos-module-rev*:

assumes $x \in \text{carrier } G \ x' \in \text{carrier } G$
and $(x' \oplus \ominus x) \in H$
shows $x' \in H +> x$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-rcos-module*:
assumes $x \in \text{carrier } G \ x' \in \text{carrier } G$
shows $(x' \in H +> x) = (x' \oplus \ominus x \in H)$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-rcos-module-minus*:
includes *ring* G
assumes *carr*: $x \in \text{carrier } G \ x' \in \text{carrier } G$
shows $(x' \in H +> x) = (x' \ominus x \in H)$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-repr-independence'*:
assumes *y*: $y \in H +> x$
and *xcarr*: $x \in \text{carrier } G$
shows $H +> x = H +> y$
 ⟨*proof*⟩

lemma (in *abelian-subgroup*) *a-repr-independenceD*:
assumes *ycarr*: $y \in \text{carrier } G$
and *repr*: $H +> x = H +> y$
shows $y \in H +> x$
 ⟨*proof*⟩

13.3 Lemmas for the Set of Right Cosets

lemma (in *abelian-subgroup*) *a-rcosets-carrier*:
 $X \in \text{a-rcosets } H \implies X \subseteq \text{carrier } G$
 ⟨*proof*⟩

13.4 Addition of Subgroups

lemma (in *abelian-monoid*) *set-add-closed*:
assumes *Acarr*: $A \subseteq \text{carrier } G$
and *Bcarr*: $B \subseteq \text{carrier } G$
shows $A <+> B \subseteq \text{carrier } G$
 ⟨*proof*⟩

lemma (in *abelian-group*) *add-additive-subgroups*:
assumes *subH*: *additive-subgroup* $H \ G$
and *subK*: *additive-subgroup* $K \ G$
shows *additive-subgroup* $(H <+> K) \ G$
 ⟨*proof*⟩

end

```

theory Ideal
imports Ring AbelCoset
begin

```

14 Ideals

14.1 General definition

```

locale ideal = additive-subgroup I R + ring R +
  assumes I-l-closed:  $\llbracket a \in I; x \in \text{carrier } R \rrbracket \implies x \otimes a \in I$ 
  and I-r-closed:  $\llbracket a \in I; x \in \text{carrier } R \rrbracket \implies a \otimes x \in I$ 

```

```

interpretation ideal  $\subseteq$  abelian-subgroup I R
<proof>

```

```

lemma (in ideal) is-ideal:
  ideal I R
<proof>

```

```

lemma idealI:
  includes ring
  assumes a-subgroup: subgroup I ( $\text{carrier} = \text{carrier } R, \text{mult} = \text{add } R, \text{one} = \text{zero } R$ )
  and I-l-closed:  $\bigwedge a x. \llbracket a \in I; x \in \text{carrier } R \rrbracket \implies x \otimes a \in I$ 
  and I-r-closed:  $\bigwedge a x. \llbracket a \in I; x \in \text{carrier } R \rrbracket \implies a \otimes x \in I$ 
  shows ideal I R
<proof>

```

14.2 Ideals Generated by a Subset of *carrier R*

```

constdefs (structure R)
  genideal :: ('a, 'b) ring-scheme  $\Rightarrow$  'a set  $\Rightarrow$  'a set (Idl1 - [80] 79)
  genideal R S  $\equiv$  Inter {I. ideal I R  $\wedge$  S  $\subseteq$  I}

```

14.3 Principal Ideals

```

locale principalideal = ideal +
  assumes generate:  $\exists i \in \text{carrier } R. I = \text{Idl } \{i\}$ 

```

```

lemma (in principalideal) is-principalideal:
  shows principalideal I R
<proof>

```

```

lemma principalidealI:
  includes ideal
  assumes generate:  $\exists i \in \text{carrier } R. I = \text{Idl } \{i\}$ 
  shows principalideal I R
<proof>

```

14.4 Maximal Ideals

locale *maximalideal* = *ideal* +
assumes *I-notcarr*: *carrier R* \neq *I*
and *I-maximal*: $\llbracket \text{ideal } J \text{ } R; I \subseteq J; J \subseteq \text{carrier } R \rrbracket \implies J = I \vee J = \text{carrier } R$

lemma (*in maximalideal*) *is-maximalideal*:
shows *maximalideal I R*
 $\langle \text{proof} \rangle$

lemma *maximalidealI*:
includes *ideal*
assumes *I-notcarr*: *carrier R* \neq *I*
and *I-maximal*: $\bigwedge J. \llbracket \text{ideal } J \text{ } R; I \subseteq J; J \subseteq \text{carrier } R \rrbracket \implies J = I \vee J = \text{carrier } R$
shows *maximalideal I R*
 $\langle \text{proof} \rangle$

14.5 Prime Ideals

locale *primeideal* = *ideal* + *cring* +
assumes *I-notcarr*: *carrier R* \neq *I*
and *I-prime*: $\llbracket a \in \text{carrier } R; b \in \text{carrier } R; a \otimes b \in I \rrbracket \implies a \in I \vee b \in I$

lemma (*in primeideal*) *is-primeideal*:
shows *primeideal I R*
 $\langle \text{proof} \rangle$

lemma *primeidealI*:
includes *ideal*
includes *cring*
assumes *I-notcarr*: *carrier R* \neq *I*
and *I-prime*: $\bigwedge a \ b. \llbracket a \in \text{carrier } R; b \in \text{carrier } R; a \otimes b \in I \rrbracket \implies a \in I \vee b \in I$
shows *primeideal I R*
 $\langle \text{proof} \rangle$

lemma *primeidealI2*:
includes *additive-subgroup I R*
includes *cring*
assumes *I-l-closed*: $\bigwedge a \ x. \llbracket a \in I; x \in \text{carrier } R \rrbracket \implies x \otimes a \in I$
and *I-r-closed*: $\bigwedge a \ x. \llbracket a \in I; x \in \text{carrier } R \rrbracket \implies a \otimes x \in I$
and *I-notcarr*: *carrier R* \neq *I*
and *I-prime*: $\bigwedge a \ b. \llbracket a \in \text{carrier } R; b \in \text{carrier } R; a \otimes b \in I \rrbracket \implies a \in I \vee b \in I$
shows *primeideal I R*
 $\langle \text{proof} \rangle$

15 Properties of Ideals

15.1 Special Ideals

lemma (in ring) zeroideal:
 shows ideal $\{0\}$ R
 ⟨proof⟩

lemma (in ring) oneideal:
 shows ideal (carrier R) R
 ⟨proof⟩

lemma (in domain) zeroprimeideal:
 shows primeideal $\{0\}$ R
 ⟨proof⟩

15.2 General Ideal Properties

lemma (in ideal) one-imp-carrier:
 assumes I -one-closed: $1 \in I$
 shows $I = \text{carrier } R$
 ⟨proof⟩

lemma (in ideal) I carr:
 assumes iI : $i \in I$
 shows $i \in \text{carrier } R$
 ⟨proof⟩

15.3 Intersection of Ideals

Intersection of two ideals The intersection of any two ideals is again an ideal in R

lemma (in ring) i -intersect:
 includes ideal I R
 includes ideal J R
 shows ideal $(I \cap J)$ R
 ⟨proof⟩

15.3.1 Intersection of a Set of Ideals

The intersection of any Number of Ideals is again an Ideal in R

lemma (in ring) i -Intersect:
 assumes S ideals: $\bigwedge I. I \in S \implies \text{ideal } I$ R
 and notempty: $S \neq \{\}$
 shows ideal (Inter S) R
 ⟨proof⟩

15.4 Addition of Ideals

lemma (in ring) *add-ideals*:
 assumes *idealI*: ideal I R
 and *idealJ*: ideal J R
 shows ideal $(I <+> J)$ R
 ⟨*proof*⟩

15.5 Ideals generated by a subset of *carrier* R

15.5.1 Generation of Ideals in General Rings

genideal generates an ideal

lemma (in ring) *genideal-ideal*:
 assumes *Scarr*: $S \subseteq \text{carrier } R$
 shows ideal $(\text{Idl } S)$ R
 ⟨*proof*⟩

lemma (in ring) *genideal-self*:
 assumes $S \subseteq \text{carrier } R$
 shows $S \subseteq \text{Idl } S$
 ⟨*proof*⟩

lemma (in ring) *genideal-self'*:
 assumes *carr*: $i \in \text{carrier } R$
 shows $i \in \text{Idl } \{i\}$
 ⟨*proof*⟩

genideal generates the minimal ideal

lemma (in ring) *genideal-minimal*:
 assumes *a*: ideal I R
 and *b*: $S \subseteq I$
 shows $\text{Idl } S \subseteq I$
 ⟨*proof*⟩

Generated ideals and subsets

lemma (in ring) *Idl-subset-ideal*:
 assumes *Iideal*: ideal I R
 and *Hcarr*: $H \subseteq \text{carrier } R$
 shows $(\text{Idl } H \subseteq I) = (H \subseteq I)$
 ⟨*proof*⟩

lemma (in ring) *subset-Idl-subset*:
 assumes *Icarr*: $I \subseteq \text{carrier } R$
 and *HI*: $H \subseteq I$
 shows $\text{Idl } H \subseteq \text{Idl } I$
 ⟨*proof*⟩

lemma (in ring) *Idl-subset-ideal'*:

assumes $acarr: a \in \text{carrier } R$ **and** $bcarr: b \in \text{carrier } R$
shows $(\text{Idl } \{a\} \subseteq \text{Idl } \{b\}) = (a \in \text{Idl } \{b\})$
 $\langle \text{proof} \rangle$

lemma (**in ring**) *genideal-zero*:
 $\text{Idl } \{0\} = \{0\}$
 $\langle \text{proof} \rangle$

lemma (**in ring**) *genideal-one*:
 $\text{Idl } \{1\} = \text{carrier } R$
 $\langle \text{proof} \rangle$

15.5.2 Generation of Principal Ideals in Commutative Rings

constdefs (**structure** R)
 $cgenideal :: ('a, 'b) \text{ monoid-scheme} \Rightarrow 'a \Rightarrow 'a \text{ set} \quad (\text{PIdl}_1 - [80] \ 79)$
 $cgenideal \ R \ a \equiv \{ x \otimes a \mid x. x \in \text{carrier } R \}$

$cgenideal$ (?) really generates an ideal

lemma (**in cring**) *cgenideal-ideal*:
assumes $acarr: a \in \text{carrier } R$
shows $\text{ideal } (\text{PIdl } a) \ R$
 $\langle \text{proof} \rangle$

lemma (**in ring**) *cgenideal-self*:
assumes $icarr: i \in \text{carrier } R$
shows $i \in \text{PIdl } i$
 $\langle \text{proof} \rangle$

$cgenideal$ is minimal

lemma (**in ring**) *cgenideal-minimal*:
includes $\text{ideal } J \ R$
assumes $aJ: a \in J$
shows $\text{PIdl } a \subseteq J$
 $\langle \text{proof} \rangle$

lemma (**in cring**) *cgenideal-eq-genideal*:
assumes $icarr: i \in \text{carrier } R$
shows $\text{PIdl } i = \text{Idl } \{i\}$
 $\langle \text{proof} \rangle$

lemma (**in cring**) *cgenideal-eq-rcos*:
 $\text{PIdl } i = \text{carrier } R \ \#> \ i$
 $\langle \text{proof} \rangle$

lemma (**in cring**) *cgenideal-is-principalideal*:
assumes $icarr: i \in \text{carrier } R$
shows $\text{principalideal } (\text{PIdl } i) \ R$
 $\langle \text{proof} \rangle$

15.6 Union of Ideals

lemma (*in ring*) *union-genideal*:
assumes *idealI*: *ideal I R*
and *idealJ*: *ideal J R*
shows $\text{Idl } (I \cup J) = I \langle + \rangle J$
<proof>

15.7 Properties of Principal Ideals

0 generates the zero ideal

lemma (*in ring*) *zero-genideal*:
shows $\text{Idl } \{0\} = \{0\}$
<proof>

1 generates the unit ideal

lemma (*in ring*) *one-genideal*:
shows $\text{Idl } \{1\} = \text{carrier } R$
<proof>

The zero ideal is a principal ideal

corollary (*in ring*) *zeropideal*:
shows *principalideal* $\{0\} R$
<proof>

The unit ideal is a principal ideal

corollary (*in ring*) *onepideal*:
shows *principalideal* (*carrier R*) *R*
<proof>

Every principal ideal is a right coset of the carrier

lemma (*in principalideal*) *rcos-generate*:
includes *cring*
shows $\exists x \in I. I = \text{carrier } R \#> x$
<proof>

15.8 Prime Ideals

lemma (*in ideal*) *primeidealCD*:
includes *cring*
assumes *notprime*: $\neg \text{primeideal } I R$
shows $\text{carrier } R = I \vee (\exists a b. a \in \text{carrier } R \wedge b \in \text{carrier } R \wedge a \otimes b \in I \wedge a \notin I \wedge b \notin I)$
<proof>

lemma (*in ideal*) *primeidealCE*:
includes *cring*
assumes *notprime*: $\neg \text{primeideal } I R$

obtains *carrier* $R = I$
 | $\exists a b. a \in \text{carrier } R \wedge b \in \text{carrier } R \wedge a \otimes b \in I \wedge a \notin I \wedge b \notin I$
<proof>

If $\{0\}$ is a prime ideal of a commutative ring, the ring is a domain

lemma (in *cring*) *zeroprimeideal-domainI*:

assumes *pi*: *primeideal* $\{0\}$ R

shows *domain* R

<proof>

corollary (in *cring*) *domain-eq-zeroprimeideal*:

shows *domain* $R = \text{primeideal } \{0\} R$

<proof>

15.9 Maximal Ideals

lemma (in *ideal*) *helper-I-closed*:

assumes *carr*: $a \in \text{carrier } R \ x \in \text{carrier } R \ y \in \text{carrier } R$

and *axI*: $a \otimes x \in I$

shows $a \otimes (x \otimes y) \in I$

<proof>

lemma (in *ideal*) *helper-max-prime*:

includes *cring*

assumes *acarr*: $a \in \text{carrier } R$

shows *ideal* $\{x \in \text{carrier } R. a \otimes x \in I\} R$

<proof>

In a cring every maximal ideal is prime

lemma (in *cring*) *maximalideal-is-prime*:

includes *maximalideal*

shows *primeideal* $I R$

<proof>

15.10 Derived Theorems Involving Ideals

— A non-zero cring that has only the two trivial ideals is a field

lemma (in *cring*) *trivialideals-fieldI*:

assumes *carrnzero*: $\text{carrier } R \neq \{0\}$

and *haveideals*: $\{I. \text{ideal } I R\} = \{\{0\}, \text{carrier } R\}$

shows *field* R

<proof>

lemma (in *field*) *all-ideals*:

shows $\{I. \text{ideal } I R\} = \{\{0\}, \text{carrier } R\}$

<proof>

lemma (in *cring*) *trivialideals-eq-field*:

assumes *carrnzero*: $\text{carrier } R \neq \{0\}$

shows $(\{I. \text{ideal } I R\} = \{\{0\}, \text{carrier } R\}) = \text{field } R$

<proof>

Like zeroprimeideal for domains

lemma (in *field*) *zeromaximalideal*:
maximalideal {**0**} *R*
<proof>

lemma (in *cring*) *zeromaximalideal-fieldI*:
assumes *zeromax*: *maximalideal* {**0**} *R*
shows *field* *R*
<proof>

lemma (in *cring*) *zeromaximalideal-eq-field*:
maximalideal {**0**} *R* = *field* *R*
<proof>

end

theory *RingHom*
imports *Ideal*
begin

16 Homomorphisms of Non-Commutative Rings

Lifting existing lemmas in a *ring-hom-ring* locale

locale *ring-hom-ring* = *ring* *R* + *ring* *S* + *var* *h* +
assumes *homh*: *h* ∈ *ring-hom* *R* *S*
notes *hom-mult* [*simp*] = *ring-hom-mult* [*OF homh*]
and *hom-one* [*simp*] = *ring-hom-one* [*OF homh*]

interpretation *ring-hom-cring* ⊆ *ring-hom-ring*
<proof>

interpretation *ring-hom-ring* ⊆ *abelian-group-hom* *R* *S*
<proof>

lemma (in *ring-hom-ring*) *is-ring-hom-ring*:
includes *struct* *R* + *struct* *S*
shows *ring-hom-ring* *R* *S* *h*
<proof>

lemma *ring-hom-ringI*:
includes *ring* *R* + *ring* *S*
assumes
hom-closed: !!*x*. *x* ∈ *carrier* *R* ==> *h* *x* ∈ *carrier* *S*

and compatible-mult: $\llbracket x : \text{carrier } R; y : \text{carrier } R \rrbracket \implies h (x \otimes y)$
 $= h x \otimes_S h y$
and compatible-add: $\llbracket x : \text{carrier } R; y : \text{carrier } R \rrbracket \implies h (x \oplus y) =$
 $h x \oplus_S h y$
and compatible-one: $h \mathbf{1} = \mathbf{1}_S$
shows *ring-hom-ring* $R S h$
 $\langle \text{proof} \rangle$

lemma *ring-hom-ringI2:*
includes *ring* $R + \text{ring } S$
assumes $h: h \in \text{ring-hom } R S$
shows *ring-hom-ring* $R S h$
 $\langle \text{proof} \rangle$

lemma *ring-hom-ringI3:*
includes *abelian-group-hom* $R S + \text{ring } R + \text{ring } S$
assumes *compatible-mult:* $\llbracket x : \text{carrier } R; y : \text{carrier } R \rrbracket \implies h (x \otimes y)$
 $= h x \otimes_S h y$
and compatible-one: $h \mathbf{1} = \mathbf{1}_S$
shows *ring-hom-ring* $R S h$
 $\langle \text{proof} \rangle$

lemma *ring-hom-cringI:*
includes *ring-hom-ring* $R S h + \text{cring } R + \text{cring } S$
shows *ring-hom-cring* $R S h$
 $\langle \text{proof} \rangle$

16.1 The kernel of a ring homomorphism

— the kernel of a ring homomorphism is an ideal

lemma (**in** *ring-hom-ring*) *kernel-is-ideal:*
shows *ideal* (*a-kernel* $R S h$) R
 $\langle \text{proof} \rangle$

Elements of the kernel are mapped to zero

lemma (**in** *abelian-group-hom*) *kernel-zero* [*simp*]:
 $i \in \text{a-kernel } R S h \implies h i = \mathbf{0}_S$
 $\langle \text{proof} \rangle$

16.2 Cosets

Cosets of the kernel correspond to the elements of the image of the homomorphism

lemma (**in** *ring-hom-ring*) *rcos-imp-homeq:*
assumes *acarr:* $a \in \text{carrier } R$
and *xcos:* $x \in \text{a-kernel } R S h + \triangleright a$
shows $h x = h a$
 $\langle \text{proof} \rangle$

```

lemma (in ring-hom-ring) homeq-imp-rcos:
  assumes acarr:  $a \in \text{carrier } R$ 
    and xcarr:  $x \in \text{carrier } R$ 
    and hx:  $h x = h a$ 
  shows  $x \in a\text{-kernel } R \ S \ h \ +> a$ 
  <proof>

corollary (in ring-hom-ring) rcos-eq-homeq:
  assumes acarr:  $a \in \text{carrier } R$ 
  shows  $(a\text{-kernel } R \ S \ h) \ +> a = \{x \in \text{carrier } R. h x = h a\}$ 
  <proof>

end

```

17 QuotRing: Quotient Rings

```

theory QuotRing
imports RingHom
begin

```

17.1 Multiplication on Cosets

```

constdefs (structure R)
  rcoset-mult :: [ $'a, -$ ] ring-scheme,  $'a \text{ set}, 'a \text{ set}, 'a \text{ set}$ ]  $\Rightarrow 'a \text{ set}$ 
    ([ $\text{mod } -$ ] -  $\otimes_1$  - [ $81, 81, 81$ ] 80)
  rcoset-mult R I A B  $\equiv \bigcup_{a \in A}. \bigcup_{b \in B}. I \ +> (a \otimes b)$ 

```

rcoset-mult fulfils the properties required by congruences

```

lemma (in ideal) rcoset-mult-add:
  [ $x \in \text{carrier } R; y \in \text{carrier } R$ ]  $\Longrightarrow [\text{mod } I:] (I \ +> x) \otimes (I \ +> y) = I \ +> (x$ 
 $\otimes y)$ 
  <proof>

```

17.2 Quotient Ring Definition

```

constdefs (structure R)
  FactRing :: [ $'a, 'b$ ] ring-scheme,  $'a \text{ set}$ ]  $\Rightarrow ('a \text{ set}) \text{ ring}$ 
    (infixl Quot 65)
  FactRing R I  $\equiv$ 
    ( $\text{carrier} = a\text{-rcosets } I, \text{mult} = \text{rcoset-mult } R \ I, \text{one} = (I \ +> \mathbf{1}), \text{zero} = I, \text{add}$ 
 $= \text{set-add } R)$ 

```

17.3 Factorization over General Ideals

The quotient is a ring

```

lemma (in ideal) quotient-is-ring:
  shows ring (R Quot I)

```

<proof>

This is a ring homomorphism

lemma (in *ideal*) *rcos-ring-hom*:
(op +> I) ∈ ring-hom R (R Quot I)
<proof>

lemma (in *ideal*) *rcos-ring-hom-ring*:
ring-hom-ring R (R Quot I) (op +> I)
<proof>

The quotient of a cring is also commutative

lemma (in *ideal*) *quotient-is-cring*:
includes *cring*
shows *cring (R Quot I)*
<proof>

Cosets as a ring homomorphism on crings

lemma (in *ideal*) *rcos-ring-hom-cring*:
includes *cring*
shows *ring-hom-cring R (R Quot I) (op +> I)*
<proof>

17.4 Factorization over Prime Ideals

The quotient ring generated by a prime ideal is a domain

lemma (in *primeideal*) *quotient-is-domain*:
shows *domain (R Quot I)*
<proof>

Generating right cosets of a prime ideal is a homomorphism on commutative rings

lemma (in *primeideal*) *rcos-ring-hom-cring*:
shows *ring-hom-cring R (R Quot I) (op +> I)*
<proof>

17.5 Factorization over Maximal Ideals

In a commutative ring, the quotient ring over a maximal ideal is a field. The proof follows “W. Adkins, S. Weintraub: Algebra – An Approach via Module Theory”

lemma (in *maximalideal*) *quotient-is-field*:
includes *cring*
shows *field (R Quot I)*
<proof>

end

```

theory IntRing
imports QuotRing IntDef
begin

```

18 The Ring of Integers

18.1 Some properties of *int*

```

lemma dvds-imp-abseq:
   $\llbracket l \text{ dvd } k; k \text{ dvd } l \rrbracket \implies \text{abs } l = \text{abs } (k::\text{int})$ 
  <proof>

```

```

lemma abseq-imp-dvd:
  assumes a-lk:  $\text{abs } l = \text{abs } (k::\text{int})$ 
  shows  $l \text{ dvd } k$ 
  <proof>

```

```

lemma dvds-eq-abseq:
   $(l \text{ dvd } k \wedge k \text{ dvd } l) = (\text{abs } l = \text{abs } (k::\text{int}))$ 
  <proof>

```

18.2 The Set of Integers as Algebraic Structure

18.2.1 Definition of \mathcal{Z}

```

constdefs
  int-ring :: int ring ( $\mathcal{Z}$ )
  int-ring  $\equiv$  ( $\text{carrier} = \text{UNIV}, \text{mult} = \text{op } *, \text{one} = 1, \text{zero} = 0, \text{add} = \text{op } +$ )

```

```

lemma int-Zcarr [intro!, simp]:
   $k \in \text{carrier } \mathcal{Z}$ 
  <proof>

```

```

lemma int-is-crng:
  crng  $\mathcal{Z}$ 
  <proof>

```

18.2.2 Interpretations

Since definitions of derived operations are global, their interpretation needs to be done as early as possible — that is, with as few assumptions as possible.

```

interpretation int: monoid [ $\mathcal{Z}$ ]
  where carrier  $\mathcal{Z} = \text{UNIV}$ 
  and mult  $\mathcal{Z} \ x \ y = x * y$ 
  and one  $\mathcal{Z} = 1$ 
  and pow  $\mathcal{Z} \ x \ n = x \wedge n$ 

```

⟨proof⟩

interpretation *int*: *comm-monoid* [\mathcal{Z}]

where *finprod* $\mathcal{Z} f A = (\text{if finite } A \text{ then setprod } f A \text{ else arbitrary})$

⟨proof⟩

interpretation *int*: *abelian-monoid* [\mathcal{Z}]

where *zero* $\mathcal{Z} = 0$

and *add* $\mathcal{Z} x y = x + y$

and *finsum* $\mathcal{Z} f A = (\text{if finite } A \text{ then setsum } f A \text{ else arbitrary})$

⟨proof⟩

interpretation *int*: *abelian-group* [\mathcal{Z}]

where *a-inv* $\mathcal{Z} x = -x$

and *a-minus* $\mathcal{Z} x y = x - y$

⟨proof⟩

interpretation *int*: *domain* [\mathcal{Z}]

⟨proof⟩

Removal of occurrences of *UNIV* in interpretation result — experimental.

lemma *UNIV*:

$x \in \text{UNIV} = \text{True}$

$A \subseteq \text{UNIV} = \text{True}$

$(\text{ALL } x : \text{UNIV}. P x) = (\text{ALL } x. P x)$

$(\text{EX } x : \text{UNIV}. P x) = (\text{EX } x. P x)$

$(\text{True} \dashrightarrow Q) = Q$

$(\text{True} \implies \text{PROP } R) == \text{PROP } R$

⟨proof⟩

interpretation *int* [*unfolded UNIV*]:

partial-order [(| *carrier* = *UNIV::int set*, *le* = *op* ≤ |)]

where *carrier* (| *carrier* = *UNIV::int set*, *le* = *op* ≤ |) = *UNIV*

and *le* (| *carrier* = *UNIV::int set*, *le* = *op* ≤ |) $x y = (x \leq y)$

and *lless* (| *carrier* = *UNIV::int set*, *le* = *op* ≤ |) $x y = (x < y)$

⟨proof⟩

interpretation *int* [*unfolded UNIV*]:

lattice [(| *carrier* = *UNIV::int set*, *le* = *op* ≤ |)]

where *join* (| *carrier* = *UNIV::int set*, *le* = *op* ≤ |) $x y = \max x y$

and *meet* (| *carrier* = *UNIV::int set*, *le* = *op* ≤ |) $x y = \min x y$

⟨proof⟩

interpretation *int* [*unfolded UNIV*]:

total-order [(| *carrier* = *UNIV::int set*, *le* = *op* ≤ |)]

⟨proof⟩

18.2.3 Generated Ideals of \mathcal{Z}

lemma *int-Idl*:

$Idl_{\mathcal{Z}} \{a\} = \{x * a \mid x. True\}$
 $\langle proof \rangle$

lemma *multiples-principalideal*:

principalideal $\{x * a \mid x. True\} \mathcal{Z}$
 $\langle proof \rangle$

lemma *prime-primeideal*:

assumes *prime*: $prime (nat p)$
shows *primeideal* $(Idl_{\mathcal{Z}} \{p\}) \mathcal{Z}$
 $\langle proof \rangle$

18.2.4 Ideals and Divisibility

lemma *int-Idl-subset-ideal*:

$Idl_{\mathcal{Z}} \{k\} \subseteq Idl_{\mathcal{Z}} \{l\} = (k \in Idl_{\mathcal{Z}} \{l\})$
 $\langle proof \rangle$

lemma *Idl-subset-eq-dvd*:

$(Idl_{\mathcal{Z}} \{k\} \subseteq Idl_{\mathcal{Z}} \{l\}) = (l \text{ dvd } k)$
 $\langle proof \rangle$

lemma *dvds-eq-Idl*:

$(l \text{ dvd } k \wedge k \text{ dvd } l) = (Idl_{\mathcal{Z}} \{k\} = Idl_{\mathcal{Z}} \{l\})$
 $\langle proof \rangle$

lemma *Idl-eq-abs*:

$(Idl_{\mathcal{Z}} \{k\} = Idl_{\mathcal{Z}} \{l\}) = (abs \ l = abs \ k)$
 $\langle proof \rangle$

18.2.5 Ideals and the Modulus

constdefs

$ZMod :: int \Rightarrow int \Rightarrow int \text{ set}$
 $ZMod \ k \ r == (Idl_{\mathcal{Z}} \{k\}) +>_{\mathcal{Z}} r$

lemmas *ZMod-defs* =

ZMod-def *genideal-def*

lemma *rcos-zfact*:

assumes *kIl*: $k \in ZMod \ l \ r$
shows *EX* $x. k = x * l + r$
 $\langle proof \rangle$

lemma *ZMod-imp-zmod*:

assumes *zmods*: $ZMod \ m \ a = ZMod \ m \ b$
shows $a \text{ mod } m = b \text{ mod } m$

<proof>

lemma *ZMod-mod*:

shows $ZMod\ m\ a = ZMod\ m\ (a\ mod\ m)$

<proof>

lemma *zmod-imp-ZMod*:

assumes *modeq*: $a\ mod\ m = b\ mod\ m$

shows $ZMod\ m\ a = ZMod\ m\ b$

<proof>

corollary *ZMod-eq-mod*:

shows $(ZMod\ m\ a = ZMod\ m\ b) = (a\ mod\ m = b\ mod\ m)$

<proof>

18.2.6 Factorization

constdefs

ZFact :: *int* \Rightarrow *int set ring*

ZFact *k* == $\mathcal{Z}\ Quot\ (Idl_{\mathcal{Z}}\ \{k\})$

lemmas *ZFact-defs* = *ZFact-def FactRing-def*

lemma *ZFact-is-crimg*:

shows *crimg* (*ZFact* *k*)

<proof>

lemma *ZFact-zero*:

carrier (*ZFact* 0) = $(\bigcup a.\ \{\{a\}\})$

<proof>

lemma *ZFact-one*:

carrier (*ZFact* 1) = $\{UNIV\}$

<proof>

lemma *ZFact-prime-is-domain*:

assumes *pprime*: *prime* (*nat* *p*)

shows *domain* (*ZFact* *p*)

<proof>

end

References

- [1] C. Ballarin. *Computer Algebra and Theorem Proving*. PhD thesis, University of Cambridge, 1999. <http://www4.in.tum.de/~ballarin/publications.html>.

- [2] N. Jacobson. *Basic Algebra I*. Freeman, 1985.
- [3] F. Kammüller and L. C. Paulson. A formal proof of sylow's theorem: An experiment in abstract algebra with Isabelle HOL. *J. Automated Reasoning*, (23):235–264, 1999.